

PROPERTIES OF EXTREMAL SEQUENCES FOR THE BELLMAN FUNCTION OF THE DYADIC MAXIMAL OPERATOR

ELEFTHERIOS N. NIKOLIDAKIS

ABSTRACT. We prove a necessary condition that has every extremal sequence for the Bellman function of the dyadic maximal operator. This implies the weak- L^p uniqueness for such a sequence.

1. INTRODUCTION

The dyadic maximal operator on \mathbb{R}^k is defined by

$$(1.1) \quad \mathcal{M}_d\phi(x) = \sup \left\{ \frac{1}{|Q|} \int_Q |\phi(u)| du : x \in Q, Q \subseteq \mathbb{R}^k \text{ is a dyadic cube} \right\}$$

for every $\phi \in L^1_{\text{loc}}(\mathbb{R}^k)$, where $|\cdot|$ is the Lebesgue measure on \mathbb{R}^k and the dyadic cubes are those formed by the grids $2^{-N}\mathbb{Z}^k$, $N = 0, 1, 2, \dots$.

It is well known that it satisfies the following weak type (1,1) inequality:

$$(1.2) \quad |\{x \in \mathbb{R}^k : \mathcal{M}_d\phi(x) \geq \lambda\}| \leq \frac{1}{\lambda} \int_{\{\mathcal{M}_d\phi \geq \lambda\}} |\phi(u)| du,$$

for every $\phi \in L^1(\mathbb{R}^k)$ and $\lambda > 0$.

From (1.2) it is easy to prove the following L^p -inequality

$$(1.3) \quad \|\mathcal{M}_d\phi\|_p \leq \frac{p}{p-1} \|\phi\|_p.$$

It is easy to see that (1.2) is best possible, while (1.3) is sharp as it can be seen in [W]. (See also [B1] and [B2] for general martingales).

A way of studying the dyadic maximal operator is to find certain refinements of the above inequalities. Concerning (1.2) refinements have been studied in [MN2], [N1] and [N2], while for (1.3) the Bellman function of two variables for $p > 1$, has been introduced by the following way:

$$(1.4) \quad T_p(f, F) = \sup \left\{ \frac{1}{|Q|} \int_Q (\mathcal{M}_d\phi)^p : \phi \geq 0, \frac{1}{|Q|} \int_Q \phi(u) du = f, \right. \\ \left. \frac{1}{|Q|} \int_Q \phi^p(u) du = F \right\}$$

where Q is a fixed dyadic cube on \mathbb{R}^k and $0 < f^p \leq F$.

2010 *Mathematics Subject Classification.* Primary 42B25; Secondary 42B99.

Key words and phrases. Bellman, dyadic, extremal, maximal.

The function given in (1.4) has been explicitly computed. Actually, this is done in a much more general setting of a non-atomic probability measure space (X, μ) where the dyadic sets are now given in a family of sets \mathcal{T} , called tree, which satisfies conditions similar to those that are satisfied by the dyadic cubes on $[0, 1]^k$.

Then the associated dyadic maximal operator $\mathcal{M}_{\mathcal{T}}$ is defined by

$$(1.5) \quad \mathcal{M}_{\mathcal{T}}\phi(x) = \sup \left\{ \frac{1}{\mu(I)} \int_I |\phi| d\mu : x \in I \in \mathcal{T} \right\},$$

where $\phi \in L^1(X, \mu)$.

Then the Bellman function (for a given $p > 1$) of two variables associated to $\mathcal{M}_{\mathcal{T}}$ is given by

$$(1.6) \quad S_p(f, F) = \sup \left\{ \int_X (\mathcal{M}_{\mathcal{T}}\phi)^p d\mu : \phi \geq 0, \int_X \phi d\mu = f, \int_X \phi^p d\mu = F \right\},$$

where $0 < f^p \leq F$.

In [M], (1.6) has been found to be $S_p(f, F) = F\omega_p(f^p/F)^p$ where $\omega_p : [0, 1] \rightarrow [1, \frac{p}{p-1}]$ is the inverse function H_p^{-1} of H_p defined on $[1, \frac{p}{p-1}]$ by $H_p(z) = -(p-1)z^p + pz^{p-1}$.

As a result the Bellman function is independent of the measure space (X, μ) and the underlying tree \mathcal{T} . Other approaches for the computation of (1.4) can be seen in [NM] and [SSV].

In this paper we study those sequences of functions: $(\phi_n)_n$, that are extremal for the Bellman function (1.6). That is $\phi_n : (X, \mu) \rightarrow \mathbb{R}^+$, $n = 1, 2, \dots$ satisfy $\int_X \phi_n d\mu = f$, $\int_X \phi_n^p d\mu = F$ and

$$(1.7) \quad \lim_n \int_X (\mathcal{M}_{\mathcal{T}}\phi_n)^p d\mu = F\omega_p(f^p/F)^p.$$

In Section 3 we prove the following

Theorem 1.1. *Let $\phi_n : (X, \mu) \rightarrow \mathbb{R}^+$ be as above. Then for every $I \in \mathcal{T}$,*

$$(1.8) \quad \lim_n \frac{1}{\mu(I)} \int_I \phi_n d\mu = f \quad \text{and} \quad \lim_n \frac{1}{\mu(I)} \int_I \phi_n^p d\mu = F.$$

Additionally:

$$\lim_n \frac{1}{\mu(I)} \int_I (\mathcal{M}_{\mathcal{T}}\phi_n)^p d\mu = F\omega_p(f^p/F)^p,$$

for every $I \in \mathcal{T}$. □

This gives as an immediate result that extremal functions do not exist for the Bellman function. Another corollary is the weak- L^p uniqueness of such a sequence in all interesting cases. In other words if $(\phi_n)_n, (\psi_n)_n$ are extremal sequences for (1.4), then $\lim_n \int_Q (\phi_n - \psi_n) h d\mu = 0$, for every $h \in L^p(Q)$, where $\frac{1}{p} + \frac{1}{q} = 1$. We need also to mention that related results in connection

with Kolmogorov's inequality have been treated in [MN1], while in [N3] it is given a characterization of such extremal sequences. More precisely it is proved there that they actually behave approximately like eigenfunctions of the dyadic maximal operator for a specific eigenvalue.

2. EXTREMAL SEQUENCES

Let (X, μ) be a non-atomic probability measure space. We give the following

Definition 2.1. A set \mathcal{T} of measurable subsets of X will be called a *tree* if the following are satisfied:

- i) $X \in \mathcal{T}$ and for every $I \in \mathcal{T}$, $\mu(I) > 0$.
- ii) For every $I \in \mathcal{T}$ there corresponds a finite or countable subset $C(I)$ of \mathcal{T} containing at least two elements such that
 - (a) the elements of $C(I)$ are disjoint subsets of I
 - (b) $I = \cup C(I)$
- iii) $\mathcal{T} = \bigcup_{m \geq 0} \mathcal{T}_{(m)}$, where $\mathcal{T}_{(0)} = \{X\}$ and

$$\mathcal{T}_{(m+1)} = \bigcup_{I \in \mathcal{T}_{(m)}} C(I).$$

- iv) The following holds $\lim_{m \rightarrow \infty} \sup_{I \in \mathcal{T}_{(m)}} \mu(I) = 0$. □

Definition 2.2. Given a tree \mathcal{T} we define the *maximal operator* associated to it as follows:

$$\mathcal{M}_{\mathcal{T}}\phi(x) = \sup \left\{ \frac{1}{\mu(I)} \int_I |\phi| d\mu : x \in I \in \mathcal{T} \right\}$$

for every $\phi \in L^1(X, \mu)$. □

From [M] we obtain the following:

Theorem 2.3. *The following holds*

$$\sup \left\{ \int_X (\mathcal{M}_{\mathcal{T}}\phi)^p d\mu : \phi \geq 0, \int \phi d\mu = f, \int_X \phi^p d\mu = F \right\} = F\omega_p(f^p/F)^p,$$

for $0 < f^p \leq F$. □

At last we give the following

Definition 2.4. Let $(\phi_n)_n$ be a sequence of non-negative measurable functions defined on X and $0 < f^p \leq F$, $p > 1$. $(\phi_n)_n$ is called (p, f, F) *extremal*, or simply *extremal* if the following hold:

$$\int_X \phi_n d\mu = f, \int_X \phi_n^p d\mu = F, \text{ for every } n = 1, 2, \dots$$

$$\lim_n \int_X (\mathcal{M}_{\mathcal{T}} \phi_n)^p d\mu = F \omega_p(f^p/F)^p.$$

3. MAIN THEOREM

Theorem 3.1. *Let $(\phi_n)_n$ be an extremal sequence. Then for every $I \in \mathcal{T}$ the following hold:*

- i) $\lim_n \frac{1}{\mu(I)} \int_I \phi_n d\mu = f$
- ii) $\lim_n \frac{1}{\mu(I)} \int_I \phi_n^p d\mu = F$
- iii) $\lim \frac{1}{\mu(I)} \int_I (\mathcal{M}_{\mathcal{T}} \phi_n)^p d\mu = F \omega_p(f^p/F)^p.$

Proof. We remind that $\mathcal{T}_{(0)} = \{X\}$ and $\mathcal{T} = \bigcup_{m \geq 0} \mathcal{T}_{(m)}$. We prove this theorem for $I \in \mathcal{T}_{(1)}$. Then inductively it holds for every $I \in \mathcal{T}_{(m)}$, $m \geq 1$.

Suppose then that $\mathcal{T}_{(1)} = \{I_k, k = 1, 2, \dots\}$ and $I = I_1$. We now set

$$f_{n,1} = \frac{1}{\mu(I_1)} \int_{I_1} \phi_n d\mu, \quad f_{n,2} = \frac{1}{\mu(X \setminus I_1)} \int_{X \setminus I_1} \phi_n d\mu,$$

$$(3.1) \quad F_{n,1} = \frac{1}{\mu(I_1)} \int_{I_1} \phi_n^p d\mu, \quad F_{n,2} = \frac{1}{\mu(X \setminus I_1)} \int_{X \setminus I_1} \phi_n^p d\mu, \quad \text{for } n = 1, 2, \dots,$$

The above sequences are obviously bounded, so passing to a subsequence we may suppose that

$$\lim_n f_{n,i} = f_i \quad \text{and} \quad \lim_n F_{n,i} = F_i, \quad \text{for } i = 1, 2.$$

For any $J \in \mathcal{T}$ define

$$\mathcal{M}_J \phi(t) = \sup \left\{ \frac{1}{\mu(K)} \int_K |\phi| d\mu : t \in K \in \mathcal{T}_J \right\}, \quad \text{for } t \in J,$$

where \mathcal{T}_J is defined by

$$\mathcal{T}_J = \{K \in \mathcal{T} : K \subseteq J\}.$$

Consider the measure space $(J, \frac{\mu(\cdot)}{\mu(J)})$, the tree \mathcal{T}_J and the associated maximal operator \mathcal{M}_J . Then using Theorem 2.3, we have that

$$(3.2) \quad \frac{1}{\mu(J)} \int_J (\mathcal{M}_J \phi)^p d\mu \leq \frac{1}{\mu(J)} \int_J \phi^p d\mu \cdot \omega_p \left(\frac{\left(\frac{1}{\mu(J)} \int_J \phi d\mu \right)^p}{\frac{1}{\mu(J)} \int_J \phi^p d\mu} \right)^p$$

for every $\phi \in L^p(J)$, where $\omega_p : [0, 1] \rightarrow [1, \frac{p}{p-1}]$ is H_p^{-1} , with

$$H_p(z) = -(p-1)z^p + pz^{p-1}, \quad z \in \left[1, \frac{p}{p-1}\right].$$

Since H_p is decreasing we conclude from (3.2) that

$$H_p \left(\left[\frac{\int_J (\mathcal{M}_J \phi)^p}{\int_J \phi^p d\mu} \right]^{1/p} \right) \geq \frac{1}{\mu(J)^{p-1}} \frac{(\int_J \phi d\mu)^p}{\int_J \phi^p d\mu},$$

which gives

$$\begin{aligned} & -(p-1) \int_J (\mathcal{M}_J \phi)^p d\mu + p \left(\int_J \phi^p d\mu \right)^{1/p} \cdot \left(\int_J (\mathcal{M}_J \phi)^p d\mu \right)^{1-\frac{1}{p}} \\ (3.3) \quad & = \frac{1}{\mu(J)^{p-1}} \left(\int_J \phi d\mu \right)^p + \delta_{\phi, J}, \end{aligned}$$

for some $\delta_{\phi, J} \geq 0$ positive constant depending on ϕ and J .

For $\phi = \phi_n$ and $J = I_i$, $i = 1, 2, \dots$ we obtain from (3.3)

$$\begin{aligned} & -(p-1) \int_{I_i} (\mathcal{M}_{I_i} \phi_n)^p d\mu + p \left(\int_{I_i} \phi_n^p d\mu \right)^{1/p} \cdot \left(\int_{I_i} (\mathcal{M}_{I_i} \phi_n)^p d\mu \right)^{1-\frac{1}{p}} \\ (3.4) \quad & = \frac{1}{\mu(I_i)^{p-1}} \left(\int_{I_i} \phi_n d\mu \right)^p + \delta_{n, i}, \text{ for every } n=1, 2, \dots \text{ and } i=1, 2, \dots \end{aligned}$$

Summing relations (3.4) for $i \geq 2$ we obtain

$$\begin{aligned} & -(p-1) \sum_{i=2}^{+\infty} \int_{I_i} (\mathcal{M}_{I_i} \phi_n)^p d\mu + p \sum_{i=2}^{+\infty} \left(\int_{I_i} \phi_n^p d\mu \right)^{1/p} \left(\int_{I_i} (\mathcal{M}_{I_i} \phi_n)^p d\mu \right)^{1-\frac{1}{p}} \\ (3.5) \quad & = \sum_{i=2}^{+\infty} \frac{1}{\mu(I_i)^{p-1}} \left(\int_{I_i} \phi_n d\mu \right)^p + \sum_{i=2}^{+\infty} \delta_{n, i}. \end{aligned}$$

In view now of Holder's inequality in its primitive form:

$$\sum_i a_i b_i \leq \left(\sum_i a_i^p \right)^{1/p} \left(\sum_i b_i^q \right)^{1/q},$$

for $a_i, b_i \geq 0$ and $q = p/(p-1)$, (3.5) gives

$$\begin{aligned} & -(p-1) A_2(n) + p \left(\int_{X \setminus I_1} \phi_n^p d\mu \right)^{1/p} \cdot [A_2(n)]^{1-\frac{1}{p}} \\ (3.6) \quad & \geq \sum_{i=2}^{+\infty} \frac{1}{\mu(I_i)^{p-1}} \left(\int_{I_i} \phi_n d\mu \right)^p + \sum_{i=2}^{+\infty} \delta_{n, i}, \text{ where} \end{aligned}$$

$$(3.7) \quad A_2(n) = \sum_{i=2}^{+\infty} \int_{I_i} (\mathcal{M}_{I_i} \phi_n)^p d\mu.$$

(In the last inequality we used the fact that $X \setminus I_1 = \bigcup_{i=2}^{+\infty} I_i$).

We use now Holder's inequality in the following form:

$$\frac{(\lambda_1 + \lambda_2 + \dots + \lambda_m)^p}{(\sigma_1 + \sigma_2 + \dots + \sigma_m)^{p-1}} \leq \frac{\lambda_1^p}{\sigma_1^{p-1}} + \frac{\lambda_2^p}{\sigma_2^{p-1}} + \dots + \frac{\lambda_m^p}{\sigma_m^{p-1}},$$

where $\sigma_i, \forall i = 1, 2, \dots$ and $\lambda_i \geq 0$, and obtain:

$$(3.8) \quad \sum_{i=2}^{+\infty} \frac{1}{\mu(I_i)^{p-1}} \left(\int_{I_i} \phi_n d\mu \right)^p \geq \frac{1}{\mu(X \setminus I_1)^{p-1}} \left(\int_{X \setminus I_1} \phi_n d\mu \right)^p = \mu(X \setminus I_1) f_{n,2}.$$

We also set

$$(3.9) \quad A_3(n) = \int_{X \setminus I_1} (\mathcal{M}_{\mathcal{T}} \phi_n)^p d\mu, \quad \text{for } n = 1, 2, \dots$$

Then by definition of \mathcal{M}_{I_i} we have that

$$(3.10) \quad A_3(n) \geq A_2(n).$$

From the above we then have that:

$$(3.11) \quad -(p-1)A_2(n) + p\mu(X \setminus I_1)^{1/p} (F_{n,2})^{1/p} [A_3(n)]^{1-\frac{1}{p}} = \mu(X \setminus I_1) (f_{n,2})^p + \delta_n^{(1)},$$

where $\delta_n^{(1)} \geq \sum_{i=2}^{+\infty} \delta_{n,i}$.

By passing to a subsequence we may suppose that $\lim_n A_3(n) = A_3$.

We will use now the following Lemma, the proof of which will be given at the end of this section.

Lemma 3.2. *If $(\phi_n)_n$ is extremal then we have that*

$$\lim_n \mu(\{\mathcal{M}_{\mathcal{T}} \phi_n = f\}) = 0.$$

From this Lemma and Definitions (3.7) and (3.9) we easily obtain that $\lim_n A_2(n) = \lim_n A_3(n) = A_3$, in view of the fact that $I_i \in \mathcal{T}_{(1)}$ for $i = 2, 3, \dots$. Then from (3.11) we conclude that

$$(3.12) \quad \begin{aligned} & -(p-1) \int_{X \setminus I_1} (\mathcal{M}_{\mathcal{T}} \phi_n)^p d\mu + p\mu(X \setminus I_1)^{1/p} (F_{n,2})^{1/p} \left(\int_{X \setminus I_1} (\mathcal{M}_{\mathcal{T}} \phi_n)^p d\mu \right)^{1-\frac{1}{p}} \\ & = \mu(X \setminus I_1) (f_{n,2})^p + \delta_n'', \end{aligned}$$

where $\delta_n'' \geq \delta_n'$, for every $n \in N$.

In the same way we have that:

$$(3.13) \quad \begin{aligned} & -(p-1) \int_{I_1} (\mathcal{M}_{\mathcal{T}} \phi_n)^p d\mu + p\mu(I_1)^{1/p} (F_{n,1})^{1/p} \cdot \left(\int_{I_1} (\mathcal{M}_{\mathcal{T}} \phi_n)^p d\mu \right)^{1-\frac{1}{p}} \\ & = \mu(I_1) (f_{n,1})^p + \varepsilon_n'', \end{aligned}$$

where ε_n'' is such that $\varepsilon_n'' \geq \delta_{n,1}$, for every $n \in N$.

Summing now (3.12) and (3.13) and using Holder's inequality in both previously mentioned forms we have that:

$$(3.14) \quad \begin{aligned} & -(p-1) \int_X (\mathcal{M}_{\mathcal{T}} \phi_n)^p d\mu + pF^{1/p} \left(\int_X (\mathcal{M}_{\mathcal{T}} \phi_n)^p d\mu \right)^{1-\frac{1}{p}} \\ & \geq \mu(I_1) (f_{n,1})^p + \mu(X \setminus I_1) (f_{n,2})^p + \delta_n'' + \varepsilon_n'' \geq f^p + \delta_n'' + \varepsilon_n'', \end{aligned}$$

which gives

$$(3.15) \quad -(p-1) \int_X (\mathcal{M}_T \phi_1)^p d\mu + pF^{1/p} \left(\int_X (\mathcal{M}_T \phi_n)^p d\mu \right)^{1-\frac{1}{p}} = f^p + \vartheta_n,$$

where $\vartheta_n \geq \delta_n'' + \varepsilon_n''$, $n = 1, 2, \dots$.

The hypothesis now for (ϕ_n) is that

$$\lim_n \int_X (\mathcal{M}_T \phi_n)^p d\mu = F\omega_p(f^p/F)^p.$$

This gives $\vartheta_n \rightarrow 0$ in (3.15) and so

$$\delta_n'' \rightarrow 0, \varepsilon_n'' \rightarrow 0.$$

As a consequence we have

$$\mu(I_1)(f_1)^p + \mu(X \setminus I_1)(f_2)^p = f^p$$

because of equality in (3.14), as $n \rightarrow +\infty$.

Since now $\mu(I_1)f_1 + \mu(X \setminus I_1)f_2 = f$ and $t \mapsto t^p$ is strictly convex on $(0, +\infty)$ we have that $f_1 = f_2 = f$.

Additionally $\delta_n'' \rightarrow 0$, so because of (3.12) and the fact that $f_2 = f$ we immediately see that

$$(3.16) \quad \lim_n \frac{1}{\mu(X \setminus I_1)} \int_{X \setminus I_1} (\mathcal{M}_T \phi_n)^p d\mu = F_2\omega_p(f^p/F_2)^p.$$

Similarly

$$(3.17) \quad \lim_n \frac{1}{\mu(I_1)} \int_{I_1} (\mathcal{M}_T \phi_n)^p d\mu = F_1\omega_p(f^p/F_1)^p.$$

Since $(\phi_n)_n$ is extremal the last two equations give

$$(3.18) \quad \mu(I_1) \cdot F_1\omega_p(f^p/F_1)^p + \mu(X \setminus I_1) \cdot F_2\omega_p(f^p/F_2)^p = F\omega(f^p/F).$$

But as we shall prove in Lemma 3.3 below the following function $t \mapsto t\omega_p(f^p/t)^p$, $t \in (f^p, +\infty)$ is strictly concave. So since $\mu(I_1)F_1 + \mu(X \setminus I_1)F_2 = F$ we have because of (3.18) that $F_1 = F_2 = F$. Then since (3.17) holds we conclude that

$$\lim_n \frac{1}{\mu(I)} \int_I (\mathcal{M}_T \phi_n)^p d\mu = F\omega_p(f^p/F)^p,$$

and Theorem 3.1 is now proved. \square

We prove now the following

Lemma 3.3. *Let $G : (1, +\infty) \rightarrow \mathbb{R}^+$ defined by $G(t) = t\omega_p(1/t)^p$. Then G is strictly concave.*

Proof. It is known from [M] that ω_p satisfies

$$\frac{d}{dx}[\omega_p(x)]^p = -\frac{1}{p-1} \frac{\omega_p(x)}{\omega_p(x)-1}, \quad x \in [0, 1].$$

So we can easily see that

$$G'(t) = \omega_p(1/t)^p + \frac{1}{p-1} \frac{1}{t} \frac{\omega_p(1/t)}{\omega_p(1/t)-1}, \quad \text{and}$$

$$G''(t) = \frac{1}{p-1} \cdot \frac{1}{t} \left(\frac{g(t)}{g(t)-1} \right)',$$

where g is defined on $(1, +\infty)$ by $g(t) = \omega_p(1/t)$. Since $g'(t) > 0, \forall t > 1$, we have that $G''(t) < 0, \forall t > 1$ and Lemma 3.3 is proved. \square

We continue now with

Proof of Lemma 3.1: Let us suppose first that ϕ_n are \mathcal{T} -simple functions that is for every n , there exists a m_n such that ϕ_n is constant on each $I \in \mathcal{T}_{(m_n)}$. As a consequence ϕ_n is \mathcal{T} -good in the sense of [M], for every n . If we look at the proof of Lemma 9 in [M] p. 324-326 we see that in all inequalities (4.20), (4.22), (4.23), (4.24) we should have equality in the limit. So as a result we must have that $\frac{1}{(\beta+1-\beta\rho_X^n)^{p-1}} - \frac{(p-1)\beta\rho_X^n}{(\beta+1)^p} \rightarrow \frac{1}{(\beta+1)^{p-1}}$, for $\beta = \omega_p(f^p/F) - 1$, where $\rho_X^n = \frac{a_X^n}{\mu(X)} = a_X^n$, where $a_X^n = \mu(\{\mathcal{M}_{\mathcal{T}}\phi_n = f\})$. But this can happen only if $a_X^n \rightarrow 0$. So the proof is completed in the case of \mathcal{T} -simple functions. As for the general case, it is not difficult to see that if $(\phi_n)_n$ is an extremal sequence of measurable functions, then we can construct a sequence of \mathcal{T} -simple functions $(\psi_n)_n$ such that $\int_X \psi_n d\mu = f$, $\int_X \psi_n^p d\mu \leq F$ and

$$\lim_n \int_X \psi_n^p d\mu = F, \quad \lim_n \int_X (\mathcal{M}_{\mathcal{T}}\psi_n)^p d\mu = F\omega_p(f^p/F)^p.$$

Additionally, we can arrange everything in such a way that $\{\mathcal{M}_{\mathcal{T}}\phi_n = f\} \subseteq \{\mathcal{M}_{\mathcal{T}}\psi_n = f\}$.

Using the same arguments as before for $(\psi_n)_n$ we can prove that $\lim_n \mu(\{\mathcal{M}_{\mathcal{T}}\psi_n = f\}) = 0$. So $\lim_n \mu(\{\mathcal{M}_{\mathcal{T}}\phi_n = f\}) = 0$ and Lemma 3.2 is proved. \square

We now give some applications of the above.

First we prove the following

Corollary 3.4. *If $0 < f^p < F$ then there do not exist extremal functions for the Bellman function $T_p(f, F)$ described in (1.4).*

Proof. Let ϕ be an extremal function for (1.4). Applying Theorem 3.1 we see that

$$\frac{1}{\mu(I)} \int_I \phi d\mu = f \quad \text{and} \quad \frac{1}{\mu(I)} \int_I \phi^p d\mu = F,$$

for every I dyadic subcube of Q .

As we can see in [G] inequality (1.2) implies that the base of dyadic sets of the tree \mathcal{T} differentiates $L^1(Q)$. That is

$$\begin{aligned} \phi(x) &= f \quad \text{a.e.} \quad \text{and} \\ \phi^p(x) &= F \quad \text{a.e.} \end{aligned}$$

This gives $f^p = F$, which is a contradiction. \square

We also prove

Corollary 3.5. *Let $T_p(f, F)$ be described by (1.4). Then if $(\phi_n)_n, (\psi_n)_n$ are extremal sequences for this function, we must have $\phi_n - \psi_n \xrightarrow{w(L^p)} 0$, as $n \rightarrow +\infty$.*

Proof. Of course we have that

$$\lim_n \frac{1}{|I|} \int_I \phi_n(u) du = \lim_n \frac{1}{|I|} \int_I \psi_n(u) du = f.$$

So $\lim_n \int_Q (\phi_n - \psi_n) \xi_I(u) du = 0$, for every dyadic subcube $I \subseteq Q$.

Since linear combinations of the characteristic functions of the dyadic subcubes of Q are dense in $L^q(Q)$ we should have that $\lim_n \int_Q (\phi_n - \psi_n) h = 0$, for every $h \in L^q(Q)$, where $q = \frac{p}{p-1}$ that is $\phi_n - \psi_n \xrightarrow{w(L^p)} 0$, as $n \rightarrow +\infty$. \square

Acknowledgements. This research has been co-financed by the European Union and Greek national funds through the Operational Program ‘‘Education and Lifelong Learning’’ of the National Strategic Reference Framework (NSRF), Arístia code: MAXBELLMAN 2760, Research code:70/3/11913.

REFERENCES

- [B1] D. L. Burkholder, *Martingales and Fourier analysis in Banach spaces*, C.I.M.E. Lectures Varenna, Como, Italy, 1985, Lecture Notes Math. **1206** (1986), 61-108.
- [B2] D. L. Burkholder, *Explorations in martingale theory and its applications*, École d’Été de Probabilités de Saint-Flour XIX-1983, Lecture Notes Math. **1464** (1991), 1-66.
- [G] De Guzman, *Real variable methods in Fourier Analysis*, North Holland (1981).
- [M] A. D. Melas, *The Bellman functions of dyadic-like maximal operators and related inequalities*. Adv. in Math. **192** (2005), 310-340.
- [MN1] A. D. Melas, E. N. Nikolidakis, *Dyadic-like maximal operators on integrable functions and Bellman functions related to Kolmogorov’s inequality*. **362**, No 3, March 2010, 1571-1597.

- [MN2] A. D. Melas, E. N. Nikolidakis, *On weak type inequalities for dyadic maximal functions*. J. Math. Anal. Appl. **348** (2008), 404-410
- [NM] E. N. Nikolidakis, A. D. Melas, *A sharp integral rearrangement inequality for the dyadic maximal operator and applications*, arxiv:1305.2521 (Submitted)
- [N1] E. N. Nikolidakis, *Optimal weak type estimates for dyadic-like maximal operators*. Ann. Acad. Scient. Fenn. Math. **38** (2013), 229-244.
- [N2] E. N. Nikolidakis, *Sharp weak type inequalities for the dyadic maximal operator*. J. Fourier Anal. and Appl. **19** (2013), 115-139.
- [N3] E. N. Nikolidakis, *Extremal sequences for the Bellman function of the dyadic maximal operator*, arxiv:1301.2898 (Submitted).
- [SSV] L. Slavin, A. Stokolos, V. Vasyunin, *Monge-Ampere equations and Bellman functions*: CR. Math. Acad. Sci. Paris, Ser I **346** (2008), 585-588.
- [W] G. Wang, *Sharp maximal inequalities for conditionally symmetric martingales and Brownian motion*. Proc. Amer. Math. Soc. **112** (1991), 579-586

DEPARTMENT OF MATHEMATICS, NATIONAL AND KAPODISTRIAN UNIVERSITY OF
ATHENS, PANEPISTIMIOYPOLIS 15784, ATHENS, GREECE

E-mail address: `lefteeris@math.uoc.gr`