



Ευρωπαϊκή Ένωση  
Ευρωπαϊκό Κοινωνικό Ταμείο



## Κατάλογος βιβλιογραφικών αναφορών


- Adhikari, B. R. & Wagreich, M. 2011. Provenance evolution of collapse graben fill in the Himalaya—The Miocene to Quaternary Thakkola-Mustang Graben (Nepal). *Sedimentary Geology*, 1-14.
- Aguirre, J. & Sánchez-Almazo, I. M. 2004. The Messinian post-evaporitic deposits of the Gafares area (Almería-Níjar basin, SE Spain). A new view of the “Lago-Mare” facies. *Sedimentary Geology*, 168, 71-95.
- Alves, T., M. & Lourenço, S. D. N. 2010. Geomorphologic features related to gravitational collapse: Submarine landsliding to lateral spreading on a Late Miocene–Quaternary slope (SE Crete, eastern Mediterranean). *Geomorphology*, 123, 13-33.
- Anapliotis, K. 1967. Contribution à la connaissance de Néogène d’Hiraklion (région Chersonisos). *Ann. Géol. Pays Hellén.*, 27, 96-109.
- Anastasakis, G. & Piper, D. J. W. 2005. Late Neogene evolution of the western South Aegean volcanic arc: sedimentary imprint of volcanicity around Milos. *Marine Geology*, 215, 135-158.
- Anastasakis, G., Piper, D. J. W., Dermitzakis, M. D. & Karakitsios, V. 2006. Upper Cenozoic stratigraphy and paleogeographic evolution of Myrtoon and adjacent basins, Aegean Sea, Greece. *Marine and Petroleum Geology*, 23, 353-369.
- Anastasakis, G. A., Dermitzakis, M. D. T. & Triantaphyllou, M. V. 1995. Stratigraphic Framework of the Gavdos island Neogene sediments. *Newsletters in Stratigraphy*, 32, 1-15.
- Andersen, N., Paul, H., Bernasconi, S. M., Mckenzie, J. A., Behrens, A., Schaeffer, P. & Albrecht, P. 2001. Large and rapid climate variability during the Messinian salinity crisis: Evidence from deuterium concentrations of individual biomarkers. *Geology*, 29, 799-802.
- Andre, J.-P., Cornee, J.-J., Saint Martin, J.-P. & Lapointe, P. 2002. Les carbonates récifaux messiniens du seuil pélagien à Lampedusa (Méditerranée centrale). *Geodiversitas*, 24, 1-15.
- Andre, J.-P., Saint Martin, S., Moissette, P., Garcia, F., Cornee, J.-J. & Ferrandini, M. 2004. An unusual Messinian succession in the Sinis Peninsula, western Sardinia, Italy. *Sedimentary Geology*, 167, 41-55.
- Angelier, J. 1979. *Neotectonique de l'arc Egeen*. PhD Thesis, D'etat. Univ. P. et M. Curie,
- Angelier, J., Lyberis, N., Le Pichon, X., Barrier, E. & Huchon, P. 1982. The tectonic development of the Hellenic arc and the sea of Crete: A synthesis. *Tectonophysics*, 86, 159-196.
- Angelone, C., Colombero, S., Esu, D., Giuntelli, P., Marcolini, F., Pavia, M., Trenkwalder, S., Van Den Hoek Ostende, L. W., Zunino, M. & Pavia, G. 2011. Moncucco Torinese, a new post-evaporitic Messinian fossiliferous site from Piedmont (NW Italy). *Neues Jahrbuch für Geologie und Paleontologie - Abhandlungen*, 259, 89-104.
- Antonarakou, A., Dermitzakis, M. D., Drinia, H., Triantaphyllou, M. V. & Tsaparas, N. 1998. Biostratigraphical and paleoecological analysis of the neogene

deposits of Kalamos island, Ionian Sea, Greece. *Bulletin of the Geological Society of Greece*, XXXII, 125-132.

- Antonarakou, A., Dermitzakis, M. D., Drinia, H., Triantaphyllou, M. V. & Tsaparas, N. 2000. Study of the Late Miocene fauna of Faneromení section, Eastern Crete. Biostratigraphical implications and cyclostratigraphical patterns. *Proceedings RCMNS, Interim-Colloquium "Mediterranean Neogene Cyclostratigraphy in marine-continental paleoenvironments"*, Geological Society of Greece, Special Publications, 9, 3-8.
- Artoni, A., Rizzini, F., Rover, M., Gennari, R., Manzi, V., Papani, G., Bernini, M., Lacombe, O., Lave, J., Roure, F. & Verges, J. 2007. Tectonic and climatic controls on sedimentation in Late Miocene Cortemaggiore Wedge-Top Basin (Northwestern Apennines, Italy). *Thrust Belts and Foreland Basins: From Fold Kinematics to Hydrocarbon Systems*. 431-456
- Bachmeyer, F., Symeonidis, N. & Zapfe, H. 1982. Die Ausgrabungen in Pikermi-Chomateri bei Athen. Eine Dokumentation. *Annalen des Naturhistorischen Museums in Wien*, 84A, 7-12.
- Bartoli, R., Catani, G., Lenardon, G. & Vinci, A. 1983. Tectonic and sedimentation of the southern Aegean Sea. *Bollettino Oceanologia Teorica ed Applicata*, 1, 319-340.
- Bassetti, M., Manzi, V., Lugli, S., Roveri, M., Longinelli, A., Lucchi, F. & Barbieri, M. 2004. Paleoenvironmental significance of Messinian post-evaporitic lacustrine carbonates in the northern Apennines, Italy. *Sedimentary Geology*, 172, 1-18.
- Bellas, S. & Frydas, D. 1994. Calcareous and siliceous phytoplankton stratigraphy from Neogene deposits of Milos island (Cyclades, Greece). *Bulletin of the Geological Society of Greece*, 30, 363-372.
- Bellas, S., Frydas, D. & Keupp, H. 2007. Late Miocene calcareous nannofossil stratigraphy and bioevents correlation: Kalidonia case section (NW Crete, Greece). *Bull. Soc. Geol. Greece*, XXXX/1, 1-13.
- Benda, L., Jonkers, H. A., Meulenkamp, J. E. & Steffens, P. 1979. Marine microfossils, sporomorphs and radiometric data from the Lower Pliocene of Ag. Thomas, Aegina, Greece. *Newsletters in Stratigraphy*, 8, 61-69.
- Benda, L., Meulenkamp, J. E. & Zachariasse, W. J. 1974. Biostratigraphic correlations in the eastern Mediterranean Neogene. Part I: Correlation between planktonic foraminiferal, uvigerinid, sporomorphal and mammal zonations of the Cretan and Italian Neogene. *Newslett. Stratigr.*, 3, 205-217.
- Biju-Duval, B. & Montadert, L. 1977. The Messinian event and recent sedimentation. In: BIJU-DUVAL, B., MONTADERT, L. (ed.) *Structural History of the Mediterranean Basins*. pp. 377-380.
- Blanc, P.-L. 2000. Of sills and straits: a quantitative assessment of the Messinian Salinity Crisis. *Deep Sea Research Part I: Oceanographic Research Papers*, 47, 1429-1460.
- Blanc, P.-L. 2006. Improved modelling of the Messinian Salinity Crisis and conceptual implications. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 238, 349-372.
- Bonneau, M., Jonkers, A. & Meulenkamp, J. E. 1984. *Geological map of Greece, Sheet Tybaki*, 1:50000. Athens: IGME.



Ευρωπαϊκή Ένωση  
Ευρωπαϊκό Κοινωνικό Ταμείο



- Butler, R. W. H., McClelland, E. & Jones, R. E. 1999. Calibrating the duration and timing of the Messinian salinity crisis in the Mediterranean: linked tectonoclimatic signals in thrustop basin of Sicily. *Journal of the Geological Society*, 156, 827–835.
- Caillère, S. & Tsoli, P. 1972. Contribution à l'histoire géologique de l'île d'Egine. *Comptes Rendus des Séances de l'Academie des Sciences, Paris*, 274, 3515–3518.
- Carnevale, G., Caputo, D. & Landini, W. 2006. Late Miocene fish otoliths from the Colombacci Formation (Northern Apennines, Italy): implications for the Messinian 'Lago-mare' event. *Geological Journal*, 41, 537-555.
- Carnevale, G., Caputo, D. & Landini, W. 2008. A leerfish (Teleostei, Carangidae) from the messinian evaporite succession of the vena del gesso basin (Romagna Apennines, Italy): Palaeogeographical and palaeoecological implications. *Bollettino Della Società Paleontologica Italiana*, 47, 196-176.
- Carnevale, G., Landini, W. & Sarti, G. 2006. Mare versus Lago-mare: marine fishes and the Mediterranean environment at the end of the Messinian Salinity Crisis. *Journal of the Geological Society*, 163, 75-80.
- Carnevale, G., Longinelli, A., Caputo, D., Barbieri, M. & Landini, W. 2008. Did the Mediterranean marine reflooding precede the Mio-Pliocene boundary? Paleontological and geochemical evidence from upper Messinian sequences of Tuscany, Italy. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 257, 81-105.
- Chaumillon, E., Mascle, J. & Hoffmann, J. 1996. Deformation of the western Mediterranean Ridge: importance of Messinian evaporitic formations. *Tectonophysics*, 263, 163-190.
- Christodoulou, G. 1961. The foraminifera of the marine Neogene of Attiki. *Geological Geophysical Research*, 7, 1-47.
- CIESM. 2008. The Messinian Salinity Crisis from Mega-deposits to Microbiology — A Consensus Report. In: BRIAND, F., ed. CIESM Workshop Monographs, 2008a Monaco. 168 pp.
- CIESM 2008. The Messinian salinity crisis. From Mega-deposits to microbiology - A consensus report. *CIESM Workshop Monographs*, 33, 168.
- Cornée, J.-J., Ferrandini, M., Saint Martin, J.-P., Münch, P., Moullade, M., Ribaud-Laurenti, A., Saint Martin, S., Roger, S. & Ferrandini, J. 2006. The late Messinian erosional surface and the subsequent reflooding in the Mediterranean: new insights from the Melilla-Nador basin (Morocco). *Paleogeography, Paleoclimatology, Paleoecology*, 230, 129-154.
- Cornée, J.-J., Léticée, J.-L., Münch, P., Quillevéré, F., Lebrun, J. F., Moissette, P., Braga, J. C., Melinte-Dobrinescu, M., De Min, L., Oudet, J. & Randrianasolo, A. in press. Sedimentology, palaeoenvironments and biostratigraphy of the Pliocene-Pleistocene carbonate platform of Grande-Terre (Guadeloupe, Lesser Antilles forearc). *Sedimentology*, 2012.
- Cornée, J.-J., Maillard, A., Conesa, G., Garcia, F., Saint Martin, J.-P., Sage, F. & Munch, P. 2008. The late Messinian erosional surface: onshore to offshore reconstruction in western Sardinia, Italy. *Sedimentary Geology*, 210, 48-60.
- Cornée, J.-J., Moissette, P., Joannin, S., Suc, J.-P., Quillevéré, F., Krijgsman, W., Hilgen, F., Münch, P., Lécuyer, C. & Desvignes, P. 2006. Tectonic and

- climatic controls of coastal sedimentation: The Late Pliocene-Early Pleistocene of northeastern Rhodes (Aegean arc, Greece). *Sedimentary Geology*, 187, 159-181.
- Cornee, J.-J., Moissette, P., Saint Martin, J.-P., Kazmer, M., Toth, E., Gorog, A., Dulai, A. & Muller, P. 2009. Marine carbonate systems in the Sarmatian (Middle Miocene) of the Central Paratethys: the Zsambek Basin of Hungary. *Sedimentology*, 56, 1728-1750.
- Cornée, J.-J., Munch, P., Quillevéré, F., Moissette, P., Vassiliev, I., Krijgsman, W., Verati, C. & Lecuyer, C. 2006. Timing of late Pliocene to middle Pleistocene tectonic events in Rhodos (Greece) inferred from  $^{40}\text{Ar}/^{39}\text{Ar}$  dating of a volcaniclastic layer. *Earth and Planetary Science Letters*, 250, 281-291.
- Cornee, J.-J., Roger, S., Munch, P., Saint Martin, J.-P., Feraud, G., Pestrea, S. & Conesa, G. 2002. Messinian events : new constraints from sedimentological investigations and new Ar39/ Ar40 ages in the Melilla-Nador basin (Morocco). *Sedimentary Geology*, 151, 127-147.
- Cornee, J.-J., Saint Martin, J.-P., Conesa, G. & Muller, J. 1994. Geometry, paleoenvironments and relative sea-level (accommodation space) changes in the Messinian Murdjadjо carbonate platform (Oran, western Algeria). Consequences. *Sedimentary Geology*, 89, 143-158.
- Cornee, J.-J., Saint Martin, J.-P., Conesa, G., Munch, P., Andre, J.-P., Saint Martin, S. & Roger, S. 2004. Correlations and sequence stratigraphic model for Messinian marine carbonate platforms of the western and central Mediterranean. *International Journal of Earth Sciences*, 93, 621-633.
- Cosentino, D., Bertini, A., Cipollari, P., Florido, F., Glioza, E., Grossi, F., Lo Mastro, S. & Sprovieri, M. 2011. Orbitally forced paleoenvironmental and paleoclimate changes in the late postevaporitic Messinian of the Central Mediterranean Basin. *Geological Society of America Bulletin*.
- De Bruijn, H., Sondaar, P. Y. & Zachariasse, W. J. 1971. Mammalia and Foraminifera from the Neogene of Kastellios Hill (Crete), a corelation of continental and marine biozones. *Koninkl. Nederl. Akad. Weten.* Amsterdam.
- Dela Pierre Francesco, F., Bernardi, E., Cavagna, S., Clari, P., Gennari, R., Irace, A., Lozar, F., Lugli, S., Manzi, V., Natalicchio, M., Roveri, M. & Violanti, D. 2011. The record of the Messinian salinity crisis in the Tertiary Piedmont Basin (NW Italy): The Alba section revisited. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 310, 238-255.
- Delrieu, B., Chaix, C., Georgiades-Dikaioula, E., Guernet, C., Koskeridou, E., Lauriat-Rage, A., Merle, D., Moissette, P. & Neraudeau, D. 1997. Paleoenvironmental and tectonosedimentary analysis of the Miocene series of Crete (Greece): evidence of a North-Cretan Back-arc basin active during the late Serravallian - earliest Tortonian times. *Neogene basins of the Med. region controls and corelation in space and time*. Catania, Italy.
- Dermitzakis, M. 1969. Geological researches of the Neogene deposits of the Ierapetra province in Crete. *Ann. Geol. Pays Hell.*, 21, 342-484.
- Dermitzakis, M. 1979. Biostratigraphic observations on the Upper Miocene deposits in Almyri Panaya section (Heraklion province, Central Crete). *Bull. Geol. Soc. of Greece*, 14, 61-69.



Ευρωπαϊκή Ένωση  
Ευρωπαϊκό Κοινωνικό Ταμείο




- Dermitzakis, M. 1979. Stratigraphy and fauna of the Upper Miocene deposits in Almyri Panayia section (Heracleion province, Central Crete). *Ann. Geol. Pays Hell.*, Tome Hors serie (III), 245-288.
- Dermitzakis, M. & Georgiades-Dikaiouli, E. 1979. Macrofauna and planktonic foraminifera of the Armenopetra section in Keratokampos - Viannou area. *Ann. Geol. Pays Hell.*, Tome Hors serie (I), 309-314.
- Dermitzakis, M. & Papanikolaou, D. 1981. Paleogeography and geodynamics of the Aegean region during the Neogene. *Ann. Geol. Pays Hell.*, Hors Serie, 245-288.
- Dermitzakis, M. D. 1978. Stratigraphy and Sedimentary History of the Miocene of Zakynthos island. *Annales géologiques des pays helléniques*, 29, 47 -186.
- Dermitzakis, M. D., Radwanska, U., Radwanski, A. & Drinia, H. 2009. Lower Pliocene (Zanclean) regressive sequence of Refina near Pikermi in Attica, Greece: A spectacular locality of mass-aggregated giant balanid cirripedes. *Hellenic Journal of Geosciences*, 44, 9 - 19.
- Dermitzakis, M. D. & Theodoridis, S. A. 1984. Biostratigraphic survey on macrofauna and microfauna and calcareous nannoplankton from the Neogene deposits of Vassiliki in East Crete. *Annales géologiques des pays helléniques*, 32, 183 -195.
- Dermitzakis, M. D. & Triantaphyllou, M. V. 1992. Contribution to the stratigraphy of Miocene sediments of Kassos Island (Southern Sporades). *N. Jb. Geol. Palaont. Mh.*, 1, 52 - 64.
- Dermitzakis, M. D. & Triantaphyllou, M. V. 1994. First order correlation of some Mediterranean upper Miocene sections. *Bulletin of the Geological Society of Greece*, XXX 317-322.
- Di Stefano, E., Cita, M. B., Spezzaferri, S. & Sprovieri, R. 1999. The Messinian-Zanclean Pissouri section (Cyprus, Eastern Mediterranean). *Memorie della Società Geologica Italiana*, 54, 133-144.
- Dietrich, V. J., Mercalli, I. & Oberhänsli, R. 1988. Dazite, High-Alumina-Basalte und Andesite als Produkte amphiboldominiierter Differentiation (Aegina und Methan, Ägäischer Inselbogen). *Schweizerische Mineralogische und Petrographische Mitteilungen*, 68, 21–39.
- Doutsos, T. & Kokkalas, S. 2001. Stress and deformation patterns in the Aegean region. *Journal of Structural Geology*, 23, 455-472.
- Drinia, H. & Antonarakou, A. 2003. Biostratigraphy, Paleoecology and use in paleoenvironmental assessment on benthic and planktonic foraminifera from the pre-evaporitic diatomaceous sediments from Gavdos island, southern Greece. In: 7ο Πανελλήνιο Συμπόσιο Ωκεανογραφίας & Αλιείας, 2003 Χερσόνησος Κρήτης, 6-9 Μαΐου. 293.
- Drinia, H., Antonarakou, A. & Kontakiotis, G. 2008. On the occurrence of Early Pliocene marine deposits in Ierapetra Basin, E. Crete. *Bulletin of Geosciences*, 83, 63-78.
- Drinia, H., Antonarakou, A., Tsaparas, N., Dermitzakis, M. D. & Kontakiotis, G. 2004. Foraminiferal record of environmental changes: pre-evaporitic diatomaceous sediments from Gavdos Island, southern Greece. *Bulletin of the Geological Society of Greece*, XXXVI/2, 782-791.



- Drinia, H., Antonarakou, A., Tsaparas, N., Dermitzakis, M. D. & Kontakiotis, G. 2004. Foraminiferal record of environmental changes: pre-evaporitic diatomaceous sediments from Gavdos Island, southern Greece. In: 10th International Congress of the Geological Society of Greece, 2004b 14-17 April, 2004, Thessaloniki. 383-384.
- Drinia, H., Antonarakou, A., Tsaparas, N. & Kontakiotis, G. 2004. Paleoecological aspects preceding the Messinian Salinity Crisis: A case from Gavdos island. In: 4th International Congress "Environment and Identity in the Mediterranean: The Messinian Salinity Crisis Revisited", 2004c 29-25 July, Corte. 33.
- Drinia, H., Antonarakou, A., Tsaparas, N. & Kontakiotis, G. 2007. Paleoenvironmental conditions preceding the Messinian Salinity Crisis: A case study from Gavdos Island. *Geobios*, 40, 251-265.
- Drinia, H., Antonarakou, A., Tsourou, T., Tzortzaki, E., Filippaki, A. & Nikolaou, C. 2010. The Messinian/Early Pliocene transition in eastern Mediterranean. New paleoenvironmental data from the Kalamaki section (Zakynthos island, Greece). In: EGU, 2010a Vienna.
- Drinia, H., Pomoni-Papaioannou, F., Triantaphyllou, M. V. & Dermitzakis, M. D. 1998. Environmental and climatological significance of the Miocene coralline algae-bryozoan facies (Apostoli Basin, Eastern Crete, Greece). In: XVI Congress of the Carpathian-Balcan Geological Association, 1998 Vienna. 142.
- Drinia, H., Pomoni-Papaioannou, F., Tsaparas, N. & Antonarakou, A. 2010. Miocene scleractinian corals of Gavdos island, southern Greece: implications for tectonic control and sea-level changes. *Bulletin of the Geological Society of Greece*, XLIII, 620-626.
- Drinia, H., Triantaphyllou, M. V. & Dermitzakis, M. D. 2001. Sedimentary facies analysis and biostratigraphical implications of the continental-marine sediments of Central-West Crete (Sellia section, Rethymnon). *Neues Jahrbuch für Geologie und Paläontologie*, 220, 295-312.
- Duermeijer, C. E., Krijgsman, W., Langereis, C. G., Meulenkamp, J. E., Triantaphyllou, M. V. & Zachariasse, W. J. 1999. A Late Pleistocene clockwise rotation phase of Zakynthos (Greece) and implications for the evolution of the western Aegean arc. *Earth and Planetary Science Letters*, 173, 315-331.
- Duermeijer, C. E., Krijgsman, W., Langereis, C. G. & Ten Veen, J. H. 1998. Post-early Messinian counterclockwise rotations on Crete: implications for Late Miocene to Recent kinematics of the southern Hellenic arc. *Tectonophysics*, 298, 177-189.
- Fassoulas, C. 1999. The structural evolution of central Crete: insight into the tectonic evolution of the south Aegean (Greece). *Journal of Geodynamics*, 27, 23-43.
- Fassoulas, C. 2001. The tectonic development of a Neogene basin at the leading edge of the active European margin: the Heraklion basin, Crete, Greece. *Journal of Geodynamics*, 31, 49-70.
- Fassoulas, C., Kiliias, A., Mountrakis, D. 1994. Postnappe stacking extension and exhumation of HP/LT rocks in the island of Crete, Greece. *Tectonics*, 13, 127-138.



Ευρωπαϊκή Ένωση  
Ευρωπαϊκό Κοινωνικό Ταμείο



- Fassoulas, C., Kilias, A. & Mountrakis, D. 1994. New data about the pre-Oligocene structural evolution of Crete (Greece)., *Bull. Soc. Geol. Greece*, 30, 67-81.
- Flecker, R., De Villiers, S. & Ellam, R. M. 2002. Modelling the effect of evaporation on the salinity– $^{87}\text{Sr}/^{86}\text{Sr}$  relationship in modern and ancient marginal-marine systems: the Mediterranean Messinian Salinity Crisis. *Earth and Planetary Science Letters*, 203, 221–233.
- Fortuin, A. R. 1977. Stratigraphy and sedimentary history of the neogene deposits in the Ierapetra region, eastern Crete. *GUA Papers of Geology*, Series 1, 164.
- Fortuin, A. R. 1978. Late Cenozoic history of Eastern Crete and implications for the geology and geodynamics of the southern Aegean sea. *Geol. Mijnb.*, 57, 451-464.
- Franco, R. L., Angela, B. M., Vinicio, M. & Marco, R. 2002. Il Messiniano trent'anni dopo: eventi connessi alla crisi di salinità nell'avanfossa appenninica. *Studi Geologici Camerti*, 1, 127-142.
- Frydas, D. 1985. Siliceous phytoplankton from a diatomite near Heraklion, Crete, Greece. *Newsletter. Stratigr.*, 14, 142-157.
- Frydas, D. 1987. Silicoflagellaten aus dem Messinium von Kreta, Griechenland. *Z. dt. Geol. Ges.*, 138, 53-75.
- Frydas, D. 1990. Stratigraphie des diatomites du Plaisancien de la Crète centrale (Grèce) à l'aide des Silicoflagellés et des nannofossiles calcaires. *Rev. Micropaléont.*, 33(2) 93-114.
- Frydas, D. 1993. Stratigraphie du Néogène de la Crète ouest (Grèce) à l'aide de Silicoflagellés et des nannofossiles calcaires. *Rev. Micropaléont.*, 36(2), 121-142.
- Frydas, D. 1994. Stratigraphie und Taxonomie von Silicoflagellaten-Vorkommen und Diatomiten des Ober-Miozän von Zentral-Kreta (Griechenland). *Berliner geowiss. Abh.*, E3, 495-505.
- Frydas, D. 1996. Upper Zanclean silicoflagellates from Milos island (Cyclades, Greece). *Journal of Nannoplankton Research*, 18, 61–67.
- Frydas, D. 1998. Upper Pliocene diatoms and silicoflagellates from section Fortessa, central Crete, Greece. *Bull. Geol. Soc. Greece*, 32(2), 93-100.
- Frydas, D. 1999. Paleoecology, stratigraphy and taxonomy of the Pliocene marine diatoms from central Crete (Greece). *Rev. Micropaléont.*, 42(4), 269-300.
- Frydas, D. 2004. Calcareous and siliceous phytoplankton stratigraphy of Neogene marine sediments in central Crete (Greece). *Revue de micropaléontologie*, 47, 87-102.
- Frydas, D. 2006. Siliceous phytoplankton biostratigraphy of the pre-evaporite Messinian diatomites in Gavdos Island, Greece. *Revue de micropaléontologie*, 49, 86-96.
- Frydas, D. & Bellas, S. 2009. Calcareous phytoplankton stratigraphy of Neogene marine sediments eastern of Heraklion, Crete, Greece. *Berliner paläobiologische Abhandlungen*, 10, 151-169.
- Frydas, D. & Keupp, H. 1992. Kieseliges und kalkiges Phytoplankton aus dem Neogen von NW- und WKreta/Griechenland. *Berliner geowiss. Abh.*, E3, 97-111.



- Frydas, D. & Keupp, H. 1996. Biostratigraphical results in Neogene deposits of NW Crete, Greece, based on calcareous nannofossils. *Berliner geowiss. Abh.*, E18, 169-189.
- Frydas, D. & Keupp, H. 2000. Biostratigraphical and paleoecological research of Lower Pliocene diatoms and silicoflagellates from Northwestern Crete, Greece. *Berliner geowiss. Abh.*, E34, 119-151.
- Frydas, D., Keupp, H. & Bellas, S. 2008. Stratigraphical investigations based on calcareous and siliceous phytoplankton assemblages from the Upper Cenozoic deposits of Messara Basin, Crete, Greece. *Z. dt. Ges. Geowiss.*, 159/3, 415-437.
- Frydas, D., Rouchy, J.-M. & Delrieu, B. 1994. Paleoecology and Stratigraphy from siliceous phytoplankton of Messinian age at central Crete, Greece. *Bull. Geol. Soc. Greece*, 30(2), 345-354.
- Fytikas, M., Innocenti, F., Kolios, N., Manetti, P. & Mazzuoli, R. 1986. The Plio-Quaternary volcanism of Saronikos area (western part of the active Aegean volcanic arc). *Annales géologiques des pays helléniques*, 33, 23-45.
- Fytrolakis, N., Peterek, A., Schroder, B. 2003. Daten zur holozanen Kustensituation bei Phaistos/Agia Triada (Messara-Ebene, Zentralkreta Griechenland) *Essener Geogr. Arb.*, 35, 57-63.
- Garcia-Castellanos, D., Estrada, F., Jiménez-Munt, I., Gorini, C., Fernández, M., Vergés, J. & De Vicente, R. 2009. Catastrophic flood of the Mediterranean after the Messinian salinity crisis. *Nature*, 462, 778-781.
- Gargani, J. & Rigollet, C. 2007. Mediterranean Sea level variations during the Messinian salinity crisis. *Geophys. Res. Lett.*, 34, L10405.
- Gaudant, J. 2008. Paleo biodiversity and paleoenvironments: The case of the teleostean fish localities of the pre-evaporitic messinian from oran and the chelif basin (Algeria). *Geodiversitas*, 30, 141-163.
- Gaudant, J., Caulet, J., Di Geronimo, I., Di Stefano, A., Fourtanier, E., Romeo, M. & Venec-Peyre, M. 1996. Sequential analysis of a new locality of fossil fishes from the diatomitic marine messinian : Masseria il salto near caltagirona (Catania province, Sicily). *Geologie Méditerranéenne*, 23, 117-147.
- Gaudant, J. & Cavallo, O. 2008. The tortonian-messinian fish faunas of piedmont (italy) and the adriatic trough: A synthesis dedicated to the memory of carlo sturani (1938-1975). *Bollettino Della Società Paleontologica Italiana*, 47, 177-189.
- Gaudant, J., Courme-Rault, M.-D., Fornaciari, E. & Fourtanier, E. 2010. The Upper Miocene fossil fish locality of Pecetto di Valenza (Piedmont, Italy): a multidisciplinary approach. *Bollettino della Società Paleontologica Italiana*, 49, 203-225.
- Gaudant, J., Fourtanier, E., Lauriat-Rage, A., Tsagaris, S., Venec-Peyre, M. & Zorn, I. 1997a. Find of a marine fish-fauna in the preevaporitic messinian of the messara (central crete, greece): Palaeoecological interpretation. *Geologie Méditerranéenne*, 24, 175-195.
- Gaudant, J., Saint-Martin, J.-P., Bessedik, M., Mansour, B., Moissette, P. & Rouchy, J.-M. 1997b. Découverte d'une frayère de poissons téléosteens dans des diatomites messiniennes du Djebel Murdjadj (environs d'Oran, Algérie). *Journal of African Earth Sciences*, 24, 511-529.



- Gaudant, J., Saint Martin, J. P., Benmoussa, A., Cornee, J. J., El Hajjaji, K. & Muller, J. 1994. Messinian fish-fauna from the periphery of the melilla-nador carbonate platform, northeastern Morocco. *Geologie Mediterraneenne*, 21, 25-35.
- Gaudant, J., Tsaparas, N., Antonarakou, A., Drinia, H., Saint-Martin, S. & Dermitzakis, M. D. 2006. A new marine fish fauna from the pre-evaporitic Messinian of Gavdos Island (Greece). *Comptes Rendus Palevol*, 5, 795-802.
- Gaudant, J., Tsaparas, N., Antonarakou, A., Drinia, H., Saint Martin, J. P. & Dermitzakis, M. D. 2006. A new marine fish fauna from the pre-evaporitic Messinian of Gavdos island (Greece). *Comptes Rendus Palevol*, 5, 795-802.
- Gelin, F. O., Sinninghe Damst  , J. S., Harrison, W. N., Maxwell, J. R. & De Leeuw, J. W. 1995. Molecular indicators for palaeoenvironmental change in a Messinian evaporitic sequence (Vena del Gesso, Italy): III. Stratigraphic changes in the molecular structure of kerogen in a single marl bed as revealed by flash pyrolysis. *Organic Geochemistry*, 23, 555.
- Gennari, R., Iaccarino, S. M., Di Stefano, A., Sturiale, G., Cipollari, P., Manzi, V., Roveri, M. & Cosentini, D. 2008. The Messinian - Zanclean boundary in the Northern Apennine. *Stratigraphy*, 5, 307-322.
- Girone, A., Nolf, D. & Cappetta, H. 2006. Pleistocene fish otoliths from the Mediterranean Basin: a synthesis. *Geobios*, 39, 651-671.
- Girone, A., Nolf, D. & Cavallo, O. 2010. Fish otoliths from the pre-evaporitic (Early Messinian) sediments of northern Italy: their stratigraphic and palaeobiogeographic significance. *Facies*, 56, 399-432.
- Glacon, G., Got, H., Rosenkrands Larsen, A., Monaco, A., Muller, C., Murat, A. 1983. Le Pliocene immerge du mole Sud-Est de Gavdos (Mediterranee orientale): implications tectoniques. *Comptes Rendus hebdomadaires des Seances de l'Academie des Sciences*, II, 363-366.
- Gleixner, G., M  gler, I. & Todd, E. D. a. R. T. W. S. 2007. Compound-Specific Hydrogen Isotope Ratios of Biomarkers: Tracing Climatic Changes in the Past. *Terrestrial Ecology*. Elsevier. 249
- Grice, K., Schouten, S., Nissenbaum, A., Charrach, J. & Sinninghe Damst  , J. S. 1998a. A remarkable paradox: Sulfurised freshwater algal (*Botryococcus braunii*) lipids in an ancient hypersaline euxinic ecosystem. *Organic Geochemistry*, 28, 195.
- Grice, K., Schouten, S., Peters, K. E. & Sinninghe Damst  , J. S. 1998b. Molecular isotopic characterisation of hydrocarbon biomarkers in Palaeocene-Eocene evaporitic, lacustrine source rocks from the Jianghan Basin, China. *Organic Geochemistry*, 29, 1745.
- Guido, A., Jacob, J. R. M., Gautret, P., Laggoun-D  arge, F., Mastandrea, A. & Russo, F. 2007. Molecular fossils and other organic markers as palaeoenvironmental indicators of the Messinian Calcare di Base Formation: normal versus stressed marine deposition (Rossano Basin, northern Calabria, Italy). *Palaeogeography, Palaeoclimatology, Palaeoecology*, 255, 265.
- Hilgen, F. J., Kuiper, K. F., Krijgsman, W., Snel, E. & Laan, E. V. D. 2007. Astronomical tuning as the basis for high resolution chronostratigraphy: the intricate history of the Messinian Salinity Crisis.



- Hjelil, E., Riedl, H., Soulakelis, N., Van Den Haute, P. & Weingartner, H. 2000. Young Neogene tectonics and relief development on the Aegean islands of Naxos, Paros and Ios. *Mitteilungen der Österreichischen Geologischen Gesellschaft*, 93, 105–127.
- Hsü, K. J., Montadert, L., Bernoulli, D., Cita, M.B., Erickson, A. 1977. History of the Mediterranean salinity crisis. *Nature*, 267 (5610), 399–403.
- Hsu, K. J., Montadert, L., Bernoulli, D., Bizon, G., Cita, M., Ericson, A., Fabricius, F., Garrison, R. E., Kidd, R. B., Melieres, F., Müller, C. & Wright, R. C. 1978. Site 378, Cretan Basin. *Initial Reports of the Deep Sea Drilling Project*, 42, 321–357.
- Hsu, K. J., Montadert, L., Bernoulli, D., Cita, M. B., Erickson, A., Garrison, R. E., Kidd, R. B., Melieres, F., Muller, C. & Wright, R. 1977. History of the Mediterranean salinity crisis. *Nature*, 267, 399-403.
- Hsu, K. J., Ryan, W. B. F. & Cita, M. B. 1973. Late Miocene Desiccation of the Mediterranean. *Nature*, 242, 240-244.
- Hurst, A. & Morton, A. C. 2001. Generic relationships in the mineral-chemical stratigraphy of turbidite sandstones. *Journal of the Geological Society*.
- Iaccarino, S. M., Bertini, A., Di Stefano, A., Ferraro, L., Gennari, R., Grossi, F., Lirer, F., Manzi, V., Menichetti, E., Lucchi, M. R., Taviani, M., Sturiale, G. & Angeletti, L. 2008. The Trave section (Monte dei Corvi, Ancona, Central Italy): an integrated paleontological study of the Messinian deposits. *Stratigraphy*, 5, 281-306.
- Jacobshagen, V. 1986. *Geologie von Griechenland*, Berlin-Stuttgart (Gerbrüder Bornträger).
- Joannin, S., Cornée, J.-J., Suc, J.-P., Moissette, P., Koskeridou, E., Lecuyer, C., Buisine, C., Kouli, K. & Ferry, S. 2007. Changes in vegetation and marine environments in the eastern Mediterranean during Early and Middle Pleistocene (Rhodes Island, Greece). *Journal of the Geological Society of London*, 164, 1119-1131.
- Jolivet, L., Brun, J.P., Gautier, P., Lallemand, S., Patriat, M. 1994. 3-D kinematics of extension in the Aegean from the early Miocene to the present, insights from the ductile crust. *Bull. Soc. Geol. Fr.*, 165, 195– 209.
- Jolivet, L., Goffe, B., Monie, P., Truffert, C., Patriat, M., Bonneau, M. 1996. Miocene detachment in Crete and exhumation P-T-t paths of high pressure metamorphic rocks. *Tectonics*, 15, 1129-1153.
- Jongsma, D., Wissman, G., Hinz, K. & Gardé, S. 1977. Seismic studies in the Cretan Sea. *Meteor Forschungs-Ergebnisse*, C27, 3-30.
- Karisiddaiah, S. M. 2000. Diverse methane concentrations in anoxic brines and underlying sediments, eastern Mediterranean Sea. *Deep Sea Research Part I: Oceanographic Research Papers*, 47, 1999.
- Kenig, F., Sinninghe Damst  , J. S., Frewin, N. L., Hayes, J. M. & De Leeuw, J. W. 1995. Molecular indicators for palaeoenvironmental change in a Messinian evaporitic sequence (Vena del Gesso, Italy). II: High-resolution variations in abundances and  $^{13}\text{C}$  contents of free and sulphur-bound carbon skeletons in a single marl bed. *Organic Geochemistry*, 23, 485.
- Keupp, H., Bellas, S. M., In Collab. With Frydas, D. & Bartholdy, J. 2000. Neogene development of the sedimentary basins of NW Crete Island, Chania

- Prefecture, South Aegean Arc System (Greece). *Berliner geowiss. Abh.*, E34, 3-117.
- Koskeridou, E. 2006. Pliocene Molluscs taxonomic diversity as a tool for the climatic-oceanographic reconstruction and stratigraphy. Two examples from the eastern Mediterranean. *Bull. Soc. Geol. Greece*, XXIX/I 80-86.
- Krijgsman, W. 1996. *Miocene magnetostratigraphy and cyclostratigraphy in the Mediterranean: extension of the astronomical time scale*. PhD Thesis, Utrecht 207
- Krijgsman, W., Fortuin, A. R., Hilgen, F. J. & Sierro, F. J. 2001. Astrochronology for the Messinian Sorbas basin (SE Spain) and orbital (precessional) forcing for evaporite cyclicity. *Sedimentary Geology*, 140, 43-60.
- Krijgsman, W., Hilgen, F. J., Langereis, C. G., Santarelli, A. & Zachariasse, W. J. 1995. Late Miocene magnetostratigraphy, biostratigraphy and cyclostratigraphy in the Mediterranean. *Earth and Planetary Science Letters*, 136, 475-494.
- Krijgsman, W., Hilgen, F. J., Langereis, C. G. & Zachariasse, W. J. 1994. The age of the Tortonian/Messinian boundary. *Earth and Planetary Science Letters*, 121, 533-547.
- Krijgsman, W., Hilgen, F. J., Raffi, I., Sierro, F. J. & Wilson, D. S. 1999. Chronology, causes and progression of the Messinian salinity crisis. *Nature*, 400, 652-655.
- Krijgsman, W., Hilgen, F.J., Raffi, I., Sierro, F.J., Wilson, D.S. 1999. Chronology, causes and progression of the Messinian salinity crisis. *Nature*, 400, 652–655.
- Le Pichon, X. & Angelier, J. 1979. The Hellenic arc and trench system: a key to the neotectonic evolution of the eastern Mediterranean area. *Tectonophysics*, 60, 1-42.
- Lourens, L., Hilgen, F., Shackleton, N. J., Laskar, J. & Wilson, D. 2004. Appendix 2. Orbital tuning calibration and conversions for the Neogene Period. In: F. GRADSTEIN, J. O., A. SMITH (ed.) *A Geologic Time Scale 2004* Cambridge University Press. 469-484
- Lugli, S., Bassetti, M. A., Manzi, V., Barbieri, M., Longinelli, A. & Roveri, M. 2007. The Messinian 'Vena del Gesso' evaporites revisited: Characterization of isotopic composition and organic matter. In: SCHREIBER, B. C., LUGLI, S. & BABEL, M. (eds.).
- Lugli, S., Manzi, V. & Roveri, M. 2008. New facies interpretation of the Messinian evaporites in the Mediterranean. *CIESM Workshop Monographs*, 33, 67-72.
- Münch, P., Cornée, J.-J., Feraud, G., Saint Martin, J.-P., Ferrandini, M., Garcia, F., Conesa, G., Moullade, M. & Roger, S. 2006. Precise  $^{40}\text{Ar}/^{39}\text{Ar}$  chronostratigraphy of the upper Messinian Melilla carbonate complex (NE Morocco): implications for the Messinian Salinity Crisis. *International Journal of Earth Sciences*, 95, 491-503.
- Manzi, R., Stefano, L., Vinicio, M. & Charlotte, S. B. 2008a. The shallow- to deep-water record of the Messinian salinity crisis: new insights from Sicily, Calabria and Apennine basins. *CIESM WORKSHOP MONOGRAPHS*, 33, 73-82.
- Manzi, V. 2001. Stratigrafia fisica, analisi sedimentologica microscopica e caratteri magnetostratigrafici dei depositi connessi all'evento evaporitico del Messiniano (F.n. Gessoso-solfifera 1.s.). *Stratigrafia Fisica, Analisi*



*Sedimentologica Microscopica E Caratteri Magnetostratigrafici Dei Depositi Connessi Alt'evento Evaporitico Del Messiniano.*

- Manzi, V., Iaccarino, S. M., Lugli, S. & Roveri, M. 2008b. Messinian salinity crisis revisited - II: New views of a vanished ocean Preface. *Stratigraphy*, 5, 225-226.
- Manzi, V., Lugli, S., Lucchi, F. R. & Roveri, M. 2005. Deep-water clastic evaporites deposition in the Messinian Adriatic foredeep (northern Apennines, Italy): Did the Mediterranean ever dry out? *Sedimentology*, 52, 875-902.
- Manzi, V., Lugli, S., Roveri, M. & Schreiber, B. C. 2009. A new facies model for the Upper Gypsum of Sicily (Italy): chronological and palaeoenvironmental constraints for the Messinian salinity crisis in the Mediterranean. *Sedimentology*, 56, 1937-1960.
- Manzi, V., Lugli, S., Roveri, M., Schreiber, B. C. & Gennari, R. 2011. The Messinian "Calcare di Base" (Sicily, Italy) revisited. *Geological Society of America Bulletin*, 123, 347-370.
- Manzi, V., Roveri, M., Gennari, R., Bertini, A., Biffi, U., Giunta, S., Iaccarino, S. M., Lanci, L., Lugli, S., Negri, A., Riva, A., Rossi, M. E. & Taviani, M. 2007. The deep-water counterpart of the Messinian lower evaporites in the Apennine foredeep: The fanantello section (Northern Apennines, Italy). *Palaeogeography Palaeoclimatology Palaeoecology*, 251, 470-499.
- Marco, R., Rocco, G., Stefano, L. & Vinicio, M. 2009. The Terminal Carbonate Complex: the record of sea-level changes during the Messinian salinity crisis. *GeoActa*, 8, 57-71.
- Markopoulou-Diakantoni, A., Ntouropi, A. & Kosta, I. 1998. The Cirripedia of the Pliocene deposits of Rafina (Attica, Greece). *Bulletin of the Geological Society of Greece*, 32, 73-83.
- Martin, L. 1987. *Structure et evolution recent de la mer Egée: rapports d'une étude par sismique reflexion*. PhD thesis, 305Cxix p.
- Mascle, J. & Martin, L. 1990. Shallow structure and recent evolution of the Aegean Sea: A synthesis based on continuous reflection profiles. *Marine Geology*, 94, 271-299.
- McClelland, E., Finegan, B. & Butler, R. W. H. 1996. A magnetostratigraphic study of the onset of the Mediterranean Messinian salinity crisis; Caltanissetta Basin, Sicily. *Geological Society, London, Special Publications*, 105, 205-217.
- Mcculloch, M. T. & Deckker, P. D. 1989. Sr isotope constraints on the Mediterranean environment at the end of the Messinian salinity crisis. *Nature*, 342, 62-65.
- McGovney, J. E. & Radovich, B. J. 1985. Seismic stratigraphy and facies of the Frigg Fan Complex. In: BERG, O. R., WOLVERTON, D.G. (ed.) *Seismic Stratigraphy 2: An Integrated Approach to Hydrocarbon Exploration*. 139-156
- Mckenzie, D. P. 1978. Active tectonics of the Alpine-Himalaya belt: the Aegean Sea and surrounding regions. *Geophys. J. R. Astron. Soc.*, 55, 217-254.
- Meijer, P. T. 2006. A box model of the blocked-outflow scenario for the Messinian Salinity Crisis. *Earth and Planetary Science Letters*, 248, 486-494.
- Meijer, P. T. & Krijgsman, W. 2005. A quantitative analysis of the desiccation and re-filling of the Mediterranean during the Messinian Salinity Crisis. *Earth and Planetary Science Letters*, 240, 510-520.



- Melinte-Dobrinescu, M. C., Suc, J.-P., Clauzon, G., Popescu, S.-M., Armijo, R., Meyer, B., Biltokin, D., Hagatay, M. N., Ucarkus, G., Jouannic, G., Fauquette, S. & Hakir, Z. 2009. The Messinian Salinity Crisis in the Dardanelles region: Chronostratigraphic constraints. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 278, 24-39.
- Mettos, A. 1992. *Geological and paleogeographical investigation of the continental Neogene and Quaternary formations of NE Attiki and SE Beotia*. PhD thesis, 259 p.
- Meulenkamp, J. E. 1979. The Aegean and the Messinian salinity crisis. In: Proceedings of the VI Colloquium on the Geology of the Aegean Region, 1979a Athens. 1253–1263.
- Meulenkamp, J. E. 1979. Field guide to the Neogene of Crete. *Publ. Dep. Geol. Pal.* Meulenkamp, J. E. (ed.) 1979. *Field guide to the Neogene of Crete*: Univ. Athens.
- Meulenkamp, J. E. 1985. Aspects of the Late Cenozoic evolution of the Aegean region. In: STANLEY, D. J., WEZEL, F.C. (ed.) *Geological Evolution of the Mediterranean Basin*. New York: Springer. pp. 307–321
- Meulenkamp, J. E., Driever, B. W. M., Jonkers, A., Spaak, P., Zachariasse, W. J., Van Der Zwann, G. J. 1979. Late Miocene-Pliocene climatic fluctuations and marine "cyclic" sedimentation, . *Ann. Geol. Pays Hell.*, Tome hors serie, 831-842.
- Meulenkamp, J. E., In Collaboration with Dermitzakis, M., Georgiadou-Dikeoulia, E., Jonkers, H. A. & Böger, H. 1979. Field guide to the Neogene of Crete. *Publ. Dept. Geol. Paleontol. Univ. Athens*, Ser. A, 32, 1-32.
- Meulenkamp, J. E., Van Der Zwaan, G. L. & Van Wamel, W. A. 1994. On late Miocene to recent vertical motions in the Cretan segment of the Hellenic Arc. *Tectonophysics*, 234, 53-72.
- Meulenkamp, J. E., Wortel, M. J. R., Van Wamel, W. A., Spakman, W. & Hoogerduyn-Strating, E. 1988. On the Hellenic subduction zone and the geodynamic evolution of Crete since the late middle Miocene. *Tectonophysics*, 146, 203-215.
- Moissette, P., Cornee, J.-J. & Koskeridou, E. 2010. Pleistocene rolling stones or huge bryozoan nodules in a mixed siliciclastic-carbonate environment (Rhodes, Greece). *Palaios*, 25, 22-30.
- Moissette, P., Cornée, J.-J., Mannaï-Tayech, B., Rahbi, M., Andre, J.-P. & Meon, H. 2010. The western edge of the Mediterranean Pelagian shelf: a mixed siliciclastic-carbonate Messinian ramp in northern Tunisia. *Palaeogeography Palaeoclimatology Palaeoecology*, 285, 85-103.
- Morris, A. 2000. Magnetic fabric and palaeomagnetic analyses of the Plio-Quaternary calc-alkaline series of Aegina Island, South Aegean volcanic arc, Greece. *Earth and Planetary Science Letters*, 176, 91-105.
- Morton, A. C. 1985. A new approach to provenance studies: electron microprobe analysis of detrital garnets from Middle Jurassic sandstones of the northern North Sea. *Sedimentology*.
- Morton, A. C. & Hallsworth, C. 1994. Identifying provenance-specific features of detrital heavy mineral assemblages in sandstones. *Sedimentary Geology*.
- Morton, A. C. & Hallsworth, C. R. 1999. Processes controlling the composition of heavy. *Sedimentary Geology*.



- Nechaev, V. P. & Isphording, W. C. 1993. Heavy-mineral assemblages of continental margins as indicators of plate-tectonic environments. *Journal of Sedimentary Research*.
- Negri, A. & Villa, G. 2000. Calcareous nannofossil biostratigraphy, biochronology and paleoecology at the Tortonian/Messinian boundary of the Faneromeni section (Crete). *Paleogeogr., Paleoceanogr., Paleoecol.*, 156, 195-209.
- Normark, Piper & Hiscott 1998. Sea level controls on the textural characteristics and depositional architecture of the Hueneme and associated submarine fan systems, Santa Monica Basin, California. *Sedimentology*, 45, 53-70.
- Omodeo Salé, S., Gennari, R., Lugli, S., Manzi, V. & Roveri, M. 2012. Tectonic and climatic control on the Late Messinian sedimentary evolution of the Nijar Basin (Betic Cordillera, Southern Spain). *Basin Research*.
- Otero, O., Lécuyer, C., Fourel, F., Martineau, F., Mackaye, H. T., Vignaud, P. & Brunet, M. 2011. Freshwater fish  $\delta^{18}\text{O}$  indicates a Messinian change of the precipitation regime in Central Africa. *Geology*, 39, 435-438.
- Panieri, G., Lugli, S., Manzi, V., Palinska, K. A. & Roveri, M. 2008. Microbial communities in Messinian evaporite deposits of the Vena del Gesso (northern Apennines, Italy). *Stratigraphy*, 5, 343-352.
- Panieri, G., Lugli, S., Manzi, V., Roveri, M., Schreiber, B. C. & Palinska, K. A. 2010. Ribosomal RNA gene fragments from fossilized cyanobacteria identified in primary gypsum from the late Miocene, Italy. *Geobiology*, 8, 101-111.
- Papanikolaou, D., Lykousis, V., Chronis, G. & Pavlakis, P. 1988. A comparative study of neotectonic basins across the Hellenic arc: the Messiniakos, Argolikos, Saronikos and Southern Evoikos Gulfs. *Basin Research*, 1, 167-176.
- Papanikolaou, D. & Nomikou, P. V. 1998. Neotectonic blocks and planation surfaces in Iraklion basin, Crete, Greece. *Bull. Geol. Soc. Greece*, 32(1) 231-239.
- Papanikolaou, D. & Vassilakis, E. 2000. The co-existence of E-W and NE-SW Neotectonic Structures in Messara Basin, Central Crete. *IESCA*. Izmir, Turkey.
- Papanikolaou, D. & Vassilakis, E. 2010. Thrust faults and extensional detachment faults in Cretan tectono-stratigraphy: Implications for Middle Miocene extension. *Tectonophysics*, 233-247.
- Papapetrou-Zamani, A. 1966. Contribution to the Neogene knowledge of the Heraklion area, Crete. *Ann. Geol. Pays Hellen.*, 16, 207-232.
- Papp, A., Steininger, F. F. & Georgiades-Dikeoulia, E. 1978. Biostratigraphie und Korrelation des Neogens von Trakhones südlich von Athen (Attica, Griechenland). *Annales géologiques des pays helléniques*, 29, 603–629.
- Pe-Piper, G., Piper, D. & Reynolds, P. 1983. Paleomagnetic stratigraphy and radiometric dating of the pliocene volcanic rocks of Aegina, Greece. *Bulletin of Volcanology*, 46, 1-7.
- Peckmann, J. & Thiel, V. 2004. Carbon cycling at ancient methane-seeps. *Chemical Geology*, 205, 443-467.
- Penzo, E., Gandolfi, G., Bargelloni, L., Colombo, L. & Patarnello, T. 1998. Messinian salinity crisis and the origin of freshwater lifestyle in western Mediterranean gobies. *Molecular Biology and Evolution*, 15, 1472-1480.



- Perissoratis, C. & Van Andel, T. H. 1991. Sea-level changes and tectonics in the Quaternary extensional basin of the South Evvoikos Gulf, Greece. *Terra Nova*, 3, 294-302.
- Peterek, A. & Schwarze, J. 2004. Architecture and Late Pliocene to recent evolution of outer-arc basins of the Hellenic subduction zone (south-central Crete, Greece). *Journal of Geodynamics*.
- Peters, J. M., Troelstra, S. R. & Van Harten, D. 1985. Late Neogene and quaternary vertical movements in eastern Crete and their regional significance. *J. Geol. Soc. Lond.*, 142, 501-513.
- Pierre, C., Rouchy, J. M. & Blanc-Valleron, M. M. 2002. Gas hydrate dissociation in the Lorca Basin (SE Spain) during the Mediterranean Messinian salinity crisis. *Sedimentary Geology*, 147, 247.
- Piper, D. J. W. & Perissoratis, C. 2003. Quaternary neotectonics of the South Aegean arc. *Marine Geology*, 198, 259-288.
- Piper, D. J. W., Stamatopoulos, L., Poulimenos, G., Doutsos, T. & Kontopoulos, N. 1991. Quaternary history of the Gulfs of Patras and Corinth. *Zeitschrift für Geomorphologie*, 34, 451-458.
- Pomoni-Papaioannou, F., Drinia, H. & Dermitzakis, M. D. 2003. Neogene non-tropical carbonate sedimentation in a warm temperate biogeographic province (Rethymnon Formation, Eastern Crete, Greece). *Sedimentary Geology*, 154, 147-157.
- Postma, G., Fortuin, A. R. & Van Wamel, W. A. 1993. Basin-fill patterns controlled by tectonics and climate: the Neogene "fore-arc" basins of eastern Crete as a case history. *Spec. Publs Int. Ass. Sediment.*, 20, 335-362.
- Potter, P. E., Maynard, J. B. & Depetris, P. J. 2005. *Mud & mudstones. Introduction and review*, Heidelberg, Springer-Verlag Berlin.
- Reichenbacher, B. & Kowalke, T. 2009. Neogene and present-day zoogeography of killifishes (*Aphanius* and *Aphanolebias*) in the Mediterranean and Paratethys areas. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 281, 43-56.
- Ring, U., Brachert, T. & Fassoulas, C. 2001. Middle Miocene graben development in Crete and its possible relation to large-scale detachment faults in the southern Aegean. *Terra Nova*, 13, 297-304.
- Roger, S., Munch, P., Cornee, J.-J., Saint Martin, J.-P., Feraud, G., Pestrea, S., Conesa, G. & Benmoussa, A. 2000. Ar40/Ar39 dating of the preevaporitic Messinian marine sequences of the Melilla basin (Morocco) : a proposal for some biosedimentary events as isochrons around the Alboran sea. *Earth and Planetary Science Letters*, 179, 101-113.
- Rögl, F., Bernor, R. L., Dermitzakis, M. D., Müller, C. & Stancheva, M. 1991. On the Pontian Correlation in the Aegean (Aegina Island). *Newsletters on Stratigraphy*, 24, 137-158.
- Rohling, E. J. 1999. Environmental Control on Mediterranean Salinity and  $\delta^{18}\text{O}$ . *Paleoceanography*, 14, 706-715, doi:10.1029/1999PA900042
- Rohling, E. J., Schiebel, R. & Siddall, M. 2008. Controls on Messinian Lower Evaporite cycles in the Mediterranean. *Earth and Planetary Science Letters*, 275, 165-171.
- Rouchy, J. M., Taberner, C., Blanc-Valleron, M. M., Sprovieri, R., Russell, M., Pierre, C., Di Stefano, E., Pueyo, J. J., Caruso, A., Dinarā-S-Turell, J., Gomis-

- Coll, E., Wolff, G. A., Cespuglio, G., Ditchfield, P., Pestrea, S., Combourieu-Nebout, N., Santisteban, C. & Grimalt, J. O. 1998. Sedimentary and diagenetic markers of the restriction in a marine basin: the Lorca Basin (SE Spain) during the Messinian. *Sedimentary Geology*, 121, 23.
- Roveri, M., Bertini, A., Cipollari, P., Cosentino, D., Di Stefano, A., Florindo, F., Gennari, R., Gliozzi, E., Grossi, F., Iaccarino, S., Lugli, S. & Manzi, V. 2008. "Earliest Zanclean age for the Colombacci and uppermost Di Tetto formations of the "latest Messinian" northern Apennines: New palaeoenvironmental data from the Maccarone section (Marche Province, Italy)" by Popescu et al. (2007) *Geobios* 40 (359-373). *Geobios*, 41, 669-675.
- Roveri, M., Bertini, A., Cosentino, D., Di, S. A., Gennari, R., Gliozzi, E., Grossi, F., Iaccarino, S. M., Lugli, S., Manzi, V. & Taviani, M. 2009. Messinian salinity crisis revisited-II: New views of a vanished ocean (vol 5, pg 323, 2008). *Stratigraphy*, 6, 87-87.
- Roveri, M., Bertini, A., Cosentino, D., Di Stefano, A., Gennari, R., Gliozzi, E., Grossi, F., Iaccarino, S. M., Lugli, S., Manzi, V. & Taviani, M. 2008. A high-resolution stratigraphic framework for the latest Messinian events in the Mediterranean area. *Stratigraphy*, 5, 323-342.
- Roveri, M., Gallo, A. B., Rossi, M., Gennari, R., Iaccarino, S. M., Lugli, S., Manzi, V., Negri, A., Rizzini, F. & Taviani, M. 2005. The Adriatic foreland record of Messinian events (Central Adriatic Sea, Italy). *GeoActa*, 4, 139-157.
- Roveri, M., Lugli, S., Manzi, V. & Gemmari, R. 2008. Large-scale mass wasting processes in the Messinian Ciminna Basin (Northern Sicily). *GeoActa*, 7, 81-98.
- Roveri, M., Lugli, S., Manzi, V. & Schreiber, B. C. 2008. The Messinian salinity crisis: a sequence-stratigraphic approach. *GeoActa*, 1, 117-138.
- Roveri, M., Lugli, S., Manzi, V. & Schreiber, B. C. 2008. The Messinian Sicilian stratigraphy revisited: new insights for the Messinian salinity crisis. *Terra Nova*, 20, 483-488.
- Roveri, M. & Manzi, V. 2006. The Messinian salinity crisis: Looking for a new paradigm? *Palaeogeography Palaeoclimatology Palaeoecology*, 238, 386-398.
- Roveri, M., Manzi, V., Bassetti, M. A., Merini, M. & Lucchi, F. R. 1998. Stratigraphy of the Messinian post-evaporitic stage in eastern-Romagna (northern Apennines, Italy). *Giornale di Geologia*, 60, 119-142.
- Roveri, M., Manzi, V., Gennari, R., Iaccarino, S. M. & Lugli, S. 2008. Recent advancements in the Messinian stratigraphy of Italy and their Mediterranean-scale implications. *Bollettino Della Societa Paleontologica Italiana*, 47, 71-85.
- Roveri, M., Manzi, V., Lucchi, F. & Rogledi, S. 2003. Sedimentary and tectonic evolution of the Vena del Gesso basin (Northern Apennines, Italy): Implications for the onset of the Messinian salinity crisis. *Geological Society of America Bulletin*, 115, 387-405.
- Ruggieri, G. & Sprovieri, R. 1976. Messinian salinity crisis and its paleogeographical implications. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 20, 13-21.



- Russell, M., Grimalt, J. O., Hartgers, W. A., Taberner, C. & Rouchy, J. M. 1997. Bacterial and algal markers in sedimentary organic matter deposited under natural sulphurization conditions (Lorca Basin, Murcia, Spain). *Organic Geochemistry*, 26, 605.
- Russell, M., Hartgers, W. A. & Grimalt, J. O. 2000. Identification and geochemical significance of sulphurized fatty acids in sedimentary organic matter from the Lorca Basin, SE Spain. *Geochimica et Cosmochimica Acta*, 64, 3711.
- Sage, F., Déverchère, J., Von Gronefeld, G., Gaullier, V., Gorini, C., Maillard, A. & Cornée, J.-J. 2011. Western Sardinia. In: LOFI, J. & DÉVERCHÈRE, J. E. A. (eds.) *Seismic atlas of the Messinian salinity crisis markers in the offshore Mediterranean domain*. CCGM & Mém. Soc. géol. Fr., n.s. 179. 72
- Saint Martin, J.-P. & Cornee, J.-J. 1996. The Messinian Mellila reef complex, Northeastern Rif, Morocco. In: FRANSEEN, E., ESTEBAN, M., WARD, B. & ROUCHY, J.-M. (eds.) *Models for Carbonate Stratigraphy from Miocene reef complexes of the Mediterranean Regions*. SEPM Special Publication, Concepts in Sedimentology and Paleontology Series. 227-237
- Savoye, B. & Piper, D. J. W. 1991. The Messinian event on the margin of the Mediterranean Sea in the Nice area, southern France. *Marine Geology*, 97, 279-304.
- Schaeffer, P., Reiss, C. & Albrecht, P. 1995. Geochemical study of macromolecular organic matter from sulfur-rich sediments of evaporitic origin (Messinian of Sicily) by chemical degradations. *Organic Geochemistry*, 23, 567.
- Schouten, S., Schoell, M., Rijpstra, W. I. C., Sinninghe Damstā, J. S. & De Leeuw, J. W. 1997. A molecular stable carbon isotope study of organic matter in immature Miocene Monterey sediments, Pismo basin. *Geochimica et Cosmochimica Acta*, 61, 2065.
- Schröder, B. 1986. Das postorogene Känozoikum in Griechenland/Ägäis. In: JACOBSHAGEN, V. (ed.) *Geologie von Griechenland*. Berlin: Borntraeger. pp. 209–240
- Sen, S., Valet, J.-P. & Ioakim, C. 1986. Magnetostratigraphy and biostratigraphy of the Neogene deposits of Kastellios Hill (Central Crete, Greece). *Palaeogeography, Palaeoclimatology, Palaeoecology*, 53, 321-334.
- Sgavetti, M., Pompilio, L., Roveri, M., Manzi, V., Valentino, G. M., Lugli, S., Carli, C., Amici, S., Marchese, F. & Lacava, T. 2009. Two geologic systems providing terrestrial analogues for the exploration of sulfate deposits on Mars: Initial spectral characterization. *Planetary and Space Science*, 57, 614-627.
- Sharma, T. & Clayton, R. N. 1965. Measurement of O<sub>18</sub>O<sub>16</sub> ratios of total oxygen of carbonates. *Geochimica et Cosmochimica Acta*, 29, 1347-1353.
- Sierro, F. J., Hilgen, F. J., Krijgsman, W. & Flores, J. A. 2001. The Abad composite (SE Spain): a Messinian reference section for the Mediterranean and the APTS. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 168, 141-169.
- Sinninghe Damsté, J. S., Frewin, N. L., Kenig, F. & De Leeuw, J. W. 1995. Molecular indicators for palaeoenvironmental change in a Messinian evaporitic sequence (Vena del Gesso, Italy). I: Variations in extractable organic matter of ten cyclically deposited marl beds. *Organic Geochemistry*, 23, 471.
- Snel, E., Mărunțeanu, M. & Meulenkamp, J. E. 2006. Calcareous nannofossil biostratigraphy and magnetostratigraphy of the Upper Miocene and Lower



- Pliocene of the Northern Aegean (Orphanic Gulf–Strimon Basin areas), Greece. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 238, 125-150.
- Sorbini, L. & Tirapelle Rancan, R. 1979. Messinian fossil fish of the Mediterranean. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 29, 143-154.
- Stadnitskaia, A., Bouloubassi, I., Elvert, M., Hinrichs, K. U. & Sinninghe Damst  , J. S. 2008. Extended hydroxyarchaeol, a novel lipid biomarker for anaerobic methanotrophy in cold seepage habitats. *Organic Geochemistry*, 39, 1007.
- Stamatakis, M. G. & Magganas, A. C. 1998. Thermally induced silica transformations in Pliocene diatomaceous layers from Aegina island, Greece. In: HEIN, J., OBRADOVIC, J. (ed.) *Siliceous deposits of Tethys and Pacific regions*. New York: Springer. pp. 141-150
- Stefano, L., Vinicio, M., Marco, R. & Charlotte, S. B. 2010. The Primary Lower Gypsum in the Mediterranean: A new facies interpretation for the first stage of the Messinian salinity crisis. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 297, 83-99.
- Symeonidis, N., Bachmayer, F. & Zapfe, H. 1973. Ausgrabungen in Pikermi bei Athen, Griechenland. *Annalen des Naturhistorischen Museums in Wien*, 77A, 125-132.
- Symeonidis, N. & Konstadinidis, D. 1967. Observations on the Neogenic formations of Central Crete. *Ann. G  ol. Pays Hell  n.*, 19, 657-688.
- Ten Haven, H. L., De Leeuw, J. W. & Schenck, P. A. 1985. Organic geochemical studies of a Messinian evaporitic basin, northern Apennines (Italy) I: Hydrocarbon biological markers for a hypersaline environment. *Geochimica et Cosmochimica Acta*, 49, 2181.
- Ten Veen, J. & Meijer, P. 1998. Late Miocene to Recent tectonic evolution of Crete (Greece): geological observations and model analysis. *Tectonophysics*.
- Ten Veen, J. H. 1998. *Neogene outer-arc evolution in the Cretan segment of the Hellenic arc: Tectonic, sedimentary and geodynamic reconstructions*. PhD thesis, Universiteit Utrecht, 191
- Ten Veen, J. H. & Kleinspehn, K. L. 2003. Incipient continental collision and plateboundary curvature: Late Pliocene–Holocene transtensional Hellenic forearc, Crete, Greece. *Journal of the Geological Society, London*, 160, 161-181.
- Ten Veen, J. H. & Meijer, P. T. 1998. Late Miocene to recent tectonic evolution of Crete (Greece): geological observations and model analysis. *Tectonophysics*, 298, 191-208.
- Ten Veen, J. H. & Postma, G. 1999. Neogene tectonics and basin fill patterns in the Hellenic outer-arc (Crete, Greece). *Basin Research*, 11, 223-241.
- Theodoridis, S. A. 1984. Calcareous nannofossil biozonation of the Miocene and revision of the helicoliths and discoasters. *Utrecht micropaleontological bulletins*, 32, 272p.
- Theodoropoulos, D. 1968. Stratigraphie und Tektonik des Isthmus von Megara (Griechenland). *Erlanger Geologische Abhandlungen*, 73, 3-23.
- Thomson, S. N., Stoeckert, B. & Brix, M. R. 1998. Thermochronology of the high-pressure metamorphic rocks of Crete, Greece: Implications for the speed of tectonic processes. *Geology*, 26, 259-262.



- Thorbecke, G. 1977. Die neogenen Deltaschotter von Ajia Galina/Kreta. *Berichte der Naturforschenden Gesellschaft zu Freiburg in Breisgau*, 67, 347–357.
- Tortorici, L., Caputo, R. & Monaco, C. 2010. Late Neogene to Quaternary contractional structures in Crete (Greece). *Tectonophysics*, 483, 203-213.
- Triantaphyllou, M. V. 2000. Short note on the occurrence of small Gephyrocapsa species and birefringent ceratoliths in Messinian diatomites (Gavdos Island, Southern Crete). *Journal of Nannoplankton Research*, 22, 37-40.
- Triantaphyllou, M. V. 2010. Calcareous nannofossil biostratigraphy of Langhian deposits in Lefkas (Ionian Islands). *Proceedings of the 12th International Congress, Patras, May, 2010, Bulletin of the Geological Society of Greece*, XLIII, 754-762.
- Triantaphyllou, M. V., Tsaparas, N., Stamatakis, M. G. & Dermitzakis, M. 1999. Calcareous nannofossil biostratigraphy and petrological analysis of the pre-evaporitic diatomaceous sediments from Gavdos island, Southern Greece. *Neues Jahrbuch Geologische Paläoontologische Mh.*, 3, 161-178.
- Trua, T., Manzi, V., Roveri, M. & Artoni, A. 2010. The Messinian volcaniclastic layers of the Northern Apennines: evidence for the initial phases of the Southern Tyrrhenian spreading? *Italian Journal of Geosciences*, 129, 269-279.
- Tsikouras, B., Pe-Piper, G., Piper, D. J. W. & Schaffer, M. 2011. Provenance evolution of collapse graben fill in the Himalaya—The Miocene to Quaternary Thakkola-Mustang Graben (Nepal). *Sedimentary Geology*.
- Turich, C. & Freeman, K. H. 2011. Archaeal lipids record paleosalinity in hypersaline systems. *Organic Geochemistry*, 42, 1147.
- Valente, J.-P., Laj, C., Sorel, D., Roy, S. & Valet, J.-P. 1982. Paleomagnetic results from Mio-Pliocene marine sedimentary series in Crete. *Earth and Planetary Science Letters*, 57, 159-172.
- Van Andel, T. H., Perissoratis, C. & Rondoyanni, T. 1993. Quaternary tectonics of the Argolikos Gulf and adjacent basins, Greece. *Journal of the Geological Society*, 150, 529-539.
- Van Hinsbergen, D. & Meulenkamp, J. 2006. Neogene supradetachment basin development on Crete (Greece) during exhumation of the South Aegean core complex. *Basin Research*, 103-124.
- Van Hinsbergen, D., Van Der Meer, D., Zachariasse, W. & Meulenkamp, J. 2006. Deformation of western Greece during Neogene clockwise rotation and collision with Apulia. *International Journal of Earth Sciences*, 95, 463-490.
- Van Hinsbergen, D. J. J., Krijgsman, W., Langereis, C. G., Cornee, J.-J., Duermeijer, C. E. & Van Vugt, N. 2007. Discrete Plio-Pleistocene phases of tilting and counterclockwise rotation in the southeastern Aegean arc (Rhodos ; Greece). *Journal of the Geological Society of London*, 164, 1133-1143.
- Van Hinsbergen, D. J. J., Snel, E., Garstman, S. A., Mărunțeanu, M., Langereis, C. G., Wortel, M. J. R. & Meulenkamp, J. E. 2004. Vertical motions in the Aegean volcanic arc: evidence for rapid subsidence preceding volcanic activity on Milos and Aegina. *Marine Geology*, 209, 329-345.
- Welter-Schultes, F. W. 2000. The paleogeography of late Neogene central Crete inferred from the sedimentary record combined with Albinaria land snail biogeography. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 157, 27-44.



Ευρωπαϊκή Ένωση  
Ευρωπαϊκό Κοινωνικό Ταμείο



- Zachariasse, W. J., Hinsbergen, D. J. J. V. & Fortuin, A. R. 2011. Formation and fragmentation of a lateMiocene supradetachment basin in central Crete: implications for exhumationmechanisms of high-pressure rocks in the Aegean forearc. *Basin Research*, 23, 678-701.
- Zelilidis, A., Piper, D. J. W. & Kontopoulos, N. 2002. Sedimentation and Basin Evolution of the Oligocene-Miocene Mesohellenic Basin, Greece. *AAPG Bulletin*, 86, 161-182.
- Ziegenbalg, S. B., Birgel, D., Hoffmann-Sell, L., Pierre, C., Rouchy, J. M. & Peckmann, J. 2012. Anaerobic oxidation of methane in hypersaline Messinian environments revealed by  $^{13}\text{C}$ -depleted molecular fossils. *Chemical Geology*, 292-293, 140-148.
- Ziegenbalg, S. B., Brunner, B., Rouchy, J. M., Birgel, D., Pierre, C., Böttcher, M. E., Caruso, A., Immenhauser, A. & Peckmann, J. 2010. Formation of secondary carbonates and native sulphur in sulphate-rich Messinian strata, Sicily. *Sedimentary Geology*, 227, 37-50.
- Αλεξόπουλος, Α. 1990. Γεωλογικές και υδρογεωλογικές συνθήκες της περιοχής των τοπογραφικού φύλλου "Μοχός" (Κεντροανατολική Κρήτη). Διδακτορική Διατριβή, Εθνικό και Καποδιστριακό Πανεπιστήμιο Αθηνών, 662.
- Βιδάκης, Μ., Jonkers, A. & Meulenkamp, J. E. 1994. Γεωλογικός χάρτης της Ελλάδας, Φύλλο Επάνω Αρχάναι, 1:50000. Αθήνα: ΙΓΜΕ.
- Βιδάκης, Μ. & Meulenkamp, J. E. 1988. Γεωλογικός χάρτης της Ελλάδας, Φύλλο Μοχός, 1:50000. Αθήνα: ΙΓΜΕ.
- Δαβή, Ε. & Bonneau, M. 1985. Γεωλογικός χάρτης της Ελλάδας, Φύλλο Αντισκάρι, 1:50000. Αθήνα: ΙΓΜΕ.
- Συμεωνίδης, Ν. & Κωνσταντινίδης, Δ. 1968. Παρατηρήσεις επί των νεογενών σχηματισμών της Κεντρικής Κρήτης. *Ann. Geol. Pays Hell.*, 19.
- Φυτρολάκης, Ν. 1978. Συμβολή στη γεωλογική έρευνα της Κρήτης. *Δελτίο Ελληνικής Γεωλογικής Εταιρείας*, XIII, 101-115.

### Ευχαριστίες

Η παρούσα έρευνα έχει συγχρηματοδοτηθεί από την Ευρωπαϊκή Ένωση (Ευρωπαϊκό Κοινωνικό Ταμείο - ΕΚΤ) και από εθνικούς πόρους μέσω του Επιχειρησιακού Προγράμματος «Εκπαίδευση και Δια Βίου Μάθηση» του Εθνικού Στρατηγικού Πλαισίου Αναφοράς (ΕΣΠΑ) – Ερευνητικό Χρηματοδοτούμενο Έργο: ΘΑΛΗΣ – ΕΚΠΑ “Κρίση Αλμυρότητας του Μεσσηνίου: η μεγαλύτερη περιβαλλοντική διαταραχή της Μεσογείου και οι επιπτώσεις της στον βιόκοσμο”.